

REMARKS

AMENDMENTS

IN THE CLAIMS

Applicants present amendments to claims 12 and 22 which effect changes requested by the examiner. In addition, applicants introduce new claims 24-27, and respectfully request that they be entered. These claims are limited to the present process wherein the new catalytic activity is within that IUB class already assigned to the original enzyme. Support for this new limitation is found in the specification at page 4, line 10, which introduces the IUB classification system, and from the example beginning on p.11, in which the new catalytic activity is within the IUB class assigned to the enzyme originally. It is respectfully urged that the claims are in *prima facie* condition for allowance, as the subject matter is novel and unobvious (the claims are based on extant claims which are no longer rejected over prior art), and is supported and enabled by the specification as originally filed. Entry will pose no additional burden on the examiner.

IN THE SPECIFICATION

Applicants again amend Table I on page 10 of the specification. This amendment is to introduce and/or clarify the proper trade names for the products referred to therein upon filing. Support for the amendment comes from the inherency of these names, as further set forward below. Additionally, portions of the specification

previously amended have been amended again to more clearly indicate applicants' intent.

OBJECTIONS TO THE SPECIFICATION

PARENTHETICAL STRUCTURE

In amending the parenthetical structure objected to, applicants intended that the equals sign, "=", be removed. Consistent with current amendment requirements, this sign has been struck through, like so "=". Accordingly, it should not have been retained. In each instance of the noted parenthetical structure, applicants have attempted to replace the "=" with "i.e.," or some similar term, or have simply let the parentheses serve their common grammatical function(s). It is respectfully submitted that the present amendments clarify this situation.

REDUCTION OF K_m /INCREASE OF k_{cat}

It is respectfully submitted that a reference value for K_m and/or k_{cat} is unnecessary for present purposes. In the present invention, the relative change is to be observed qualitatively, and the numerical value of the change is not important. As long as one of skill in the art can recognize that a catalytic activity was not present, using available detection methods, and subsequently recognize that the same catalytic activity *is* present, the new catalytic activity has been generated. Whether this is due specifically to a reduction in K_m or an increase in k_{cat} , and to what precise extent it is due

to either, is not necessary for one of skill in the art to know.

TRADE NAMES OF ENZYMES

The amendments to Table I on page 10 of the specification introduced the trade names of the commercially available enzymes employed in the examples beginning on page 9. Attached hereto are printed pages from various sources which indicate that the terms Amano, Lipoxyme, and Novozyme (amended herein to "Novozym," which is technically more correct) would be inherently observed by one of skill in the art when reading the information in the table as originally filed.

Applicants attach pages from the Amano Enzyme, Inc. website, which list the following products: Lipase PS "Amano," Lipase AH "Amano," Acylase "Amano," Lipase D "Amano," Lipase F-AP 15, Lipase AY "Amano," Lipase M "Amano" 10, Lipase R "Amano," and Lipase G "Amano" 50. Pages taken from the Sanger Institute website and from the German Collection of Microorganisms and Cell Cultures (DSMZ) website, also attached, indicate the equivalence between *Burkholderia cenocepacia* and *Pseudomonas cepacia*, and *Rhizopus oryzae* and strains *R. javanicus* and *R. delamar*, respectively.

Pages printed from the Novozymes A/S website indicate use of the words "Novozym" and "Lipozyme" for commercially available products manufactured by that company. The attached abstract of Yamamoto, et al. recites use of "two lipases[:] Amano PS (*Pseudomonas* sp.) and Novozym 435[®]," indicating recognized use of these

terms by skilled artisans. Likewise, the attached paper by Maugard and Legroy indicates that "Novozym[®] SP 435 (lipase from *Candida antarctica* immobilised on an acrylic resin), [and] Lipozyme[®] (lipase from *Rhizomucor miehei* immobilised on an anionic macroporous resin ...), were [obtained] from Novo Industries (Denmark)." Novo has since moved its industrial enzyme production to Novozymes A/S (see printed page from Novozymes referencing the Demerger Document).

Applicants respectfully submit that from these documents, as well as from a quick search of the relevant literature, the examiner may conclude that the introduced trade names are not new matter. To one of skill in the art, familiar with the commercially available enzymes, these additional terms would be inherently present in the table as originally filed.

Likewise, the supplied literature amply supports identification of the relevant enzymes as lipases.

REJECTIONS UNDER 35 USC §112. ¶2

FUNCTIONAL EQUIVALENTS

The entire context in which *E. coli* XL1 Red is discussed in the specification indicates that the relevant function of this strain is mutation. The necessary genetic markers mutS, mutT, and mutD5, now recited in the claims, each create deficiencies in the *E. coli* DNA replication process. Applicants feel that the specification provides ample indication of the relevant functional equivalence for purposes of the presently

claimed process. However, to assuage whatever residual uncertainty remains in the examiner's mind, applicants have indicated this function by appropriate amendment.

ABBREVIATIONS PS AND AH

As indicated above, Lipase PS and Lipase AH are art-recognized portions of trade names under which certain lipases from *Pseudomonas candida* are sold by Amano Enzyme, Inc. The name "Amano" is trademarked in the United States, and as such, should not be used in the claims. The letters PS and AH serve to indicate particular lipase preparations known in the art. From this, and from the above discussion of the relevant literature, what is meant by "*Pseudomonas cepacia* lipase PS" and "*Pseudomonas cepacia* lipase AH" would be abundantly clear to the skilled artisan.

REL A1

Attached to this amendment are copies of an abstract from S. Metzger, et al., (Characterization of the relA1 mutation and a comparison of relA1 with new relA null alleles in *Escherichia coli*, *J. Biol. Chem.*, 264(35):21146-52 (1989)) and an excerpt from a paper by M. K. B. Berlyn (Linkage Map of *Escherichia coli* K-12, Edition 10: The Traditional Map, *Microbiol. and Mol. Bio. Rev.*, 62(3):814-984 (1998), excerpted pages are 814-822, 834, 856, 866, 885, 888-9, 913, and 948). These documents indicate the level of knowledge held or available to one of skill in the art with reference to the relA1

mutation.

IMPEDING ENZYME ACTIVITY

Applicants have amended the claims to indicate that "a" new catalytic activity is to be introduced into the enzyme, thus making the inherent directionality express. It is respectfully submitted that one of skill in the art would recognize the need or desirability for identifying a specific catalytic activity prior to undertaking the presently claimed process. Once such an inherently necessary step has been performed, both the enzymatic activity which would impede detection and the proper substrate would also be identified.

No limitation is made in claims 12-23 as to the identity of the new catalytic activity to be obtained, or to its relation to the enzyme's original catalytic activity. The new catalytic activity to be obtained may very well be in addition to or in place of that catalytic activity possessed by the identified enzyme prior to performance of the process. The original activity may be maintained, but need not be, necessarily. Such questions would be answerable to one of skill in the art upon performing the particular iteration of the claimed process.

REJECTIONS UNDER 35 USC §112, ¶1

WRITTEN DESCRIPTION - FUNCTIONAL EQUIVALENTS

The examiner states that a "representative number" of species is required to

adequately describe the full scope of functional equivalents. However, "possession" of the invention need not be shown by an actual reduction to practice (see, e.g., *Lockwood v. American Airlines, Inc.*, 41 USPQ2d 1961, 1966 (Fed. Cir. 1997)). Rather, a compound must be defined by "whatever characteristics sufficiently distinguish it" (*Amgen, Inc. v. Chugai Pharmaceutical*, 18 USPQ2d 1016, 1021 (Fed. Cir. 1991)). There is no absolute requirement for description of an actual reduction to practice, so long as one of skill in the art would recognize that the disclosed element is obvious as disclosed, or may immediately envisage the element, relying on knowledge and level of skill in the art (see, e.g., *In re Deuel*, 34 USPQ2d 1210 (Fed. Cir. 1995); *Fujikawa v. Wattanasin*, 39 USPQ2d 1895 (Fed. Cir. 1996); *In re Ruschig*, 154 USPQ 118 (CCPA 1967)).

In the present case, the specification and claims indicate that functional equivalents of the present mutator strain are derivatives of *Escherichia coli* XL1 Red which possess the gene markers *relA1*, *mutS*, *mutT* and *mutD5*. Production of such derivatives of *E. coli* XL1 Red, given the knowledge and level of skill in the art, would be obvious and straightforward to one of ordinary skill therein. It is respectfully submitted that the present disclosure, coupled with the extensive resources available to the practitioner of genetic recombination, are sufficient to support the claimed range of functional derivatives for purposes of the written description requirement.

ENABLEMENT

Applicants reiterate their belief that the nature of the experimentation required for claims 12-23, though potentially arduous and substantial, would be a matter of routine for the skilled artisan. One of skill in the art would recognize how divergent from an enzyme's original substrate a particular substrate may be to ensure that the new catalytic activity can be produced in that enzyme. The field of enzymology is not so unpredictable that one of skill in the art would not be unable to understand the necessary parameters inherent in practicing the present invention. To answer the examiner's questions, one of skill in the art would apply the knowledge and understanding commonly held with regard to the individual enzyme to be mutated and/or substrate targeted.

CONCLUSION

In view of the foregoing amendments and remarks, applicants consider that the rejections of record have been obviated and respectfully solicit passage of the application to issue. Should the examiner disagree, applicants would sincerely appreciate suggestions as to language and/or subject matter which would be acceptable.

Please charge any shortage in fees due in connection with the filing of this paper, including Extension of Time fees to Deposit Account No. 11-0345. Please credit any excess fees to such deposit account.

BORNSCHEUER et al., Serial No. 09/161,680


Respectfully submitted,
KEIL & WEINKAUF

A handwritten signature in black ink, appearing to read 'David C. Liechty', with a long horizontal flourish extending to the right.

David C. Liechty
Reg. No. 48,692

1350 Connecticut Ave., N.W.
Washington, D.C. 20036
(202)659-0100

DCL/kas



CONTENTS

» HOME

Company Information

[About Enzymes](#)
[Product Information](#)
[Enzyme Applications](#)
[Quality Assurance](#)
[Enzyme Wave](#)

Product Information







Product
Form







Select

Biocatalysts

For inquiries or sample request

Examples of reaction types that are candidates for enzymes mediation

Application	Products	Form	Opt. pH	Opt. Temp	Speci
Chiral Synthesis	Lipase AS "Amano" <i>(Aspergillus niger)</i> Catalogue MSDS 	Powder	6.0	45°C	12,000-15,000 (Lipolytic activi
	Lipase M "Amano" 10 <i>(Mucor javanicus)</i> Catalogue MSDS 	Powder	7.0	40°C	≥ 10,000 u/g (Lipolytic activi
	Lipase F-AP 15 <i>(Rhizopus oryzae)</i> Catalogue MSDS 	Powder	6.5	40°C	≥ 150,000 u/g (Lipolytic activi
	Lipase G "Amano" 50 <i>(Penicillium camembertii)</i> Catalogue MSDS 	Powder	5.0	30°C	≥ 50,000 u/g (Lipolytic activi
	Lipase AYS "Amano" <i>(Candida rugosa)</i> Catalogue MSDS 	Powder	7.0	45°C	≥ 30,000 u/g (Lipolytic activi
	Lipase PS "Amano" <i>(Burkholderia cepacia)</i> Catalogue MSDS 	Powder	7.0	50°C	≥ 30,000 u/g (Lipolytic activi
	Lipase AK "Amano" 20 *	Powder	8.0	55°C	≥ 20,000 u/g

	(<i>Pseudomonas fluorescens</i>)					(Lipolytic activi
	Catalogue MSDS					
						
	Cholesterol Esterase "Amano" 2 (<i>Pseudomonas sp.</i>)	Powder lyophilized	7.0	35°C	≥ 10,000 u/g (Lipolytic activi	
	MSDS					
	[Lipase D "Amano" * (<i>Rhizopus oryzae</i>)	Powder lyophilized	7.0	50°C	≥ 2,000,000 u/ (Lipolytic activi	
	MSDS 					
	[Lipase AH "Amano" * (<i>Burkholderia cepacia</i>)	Powder lyophilized	7.0	50°C	≥ 8,000 u/g (Lipolytic activi	
	MSDS 					
LPS	Lipase PS-C "Amano" I * (<i>Burkholderia cepacia</i>)	Immobilized on ceramic particles	7.0	50°C	≥ 1,000 u/g (Esterification :	
	Catalogue MSDS					
	Assay Type					
	Lipase PS-C "Amano" II * (<i>Burkholderia cepacia</i>)	Immobilized on ceramic particles	7.0	50°C	≥ 600 u/g (Esterification :	
	Catalogue MSDS					
	Assay Type					
	Lipase PS-D "Amano" I * (<i>Burkholderia cepacia</i>)	Immobilized on diatomaceous earth	7.0	50°C	≥ 500 u/g (Esterification :	
	MSDS					
	[Lipase PS "Amano" (<i>Burkholderia cepacia</i>)	Powder	7.0	50°C	≥ 30,000 u/g (Lipolytic activi	
	Catalogue MSDS					
						
	LP "Amano" S (<i>Burkholderia cepacia</i>)	Powder lyophilized	7.0	50°C	≥ 2,200,000 u/ (Lipolytic activi	
	Catalogue MSDS					
	Type 					
Protease	Protease P "Amano" 10 (<i>Aspergillus melleus</i>)	Powder	8.0	45°C	≥ 100,000 u/g (Proteolytic act	
	Catalogue MSDS					
						
	Protease N "Amano" (<i>Bacillus subtilis</i>)	Powder	7.0	55°C	≥ 150,000 u/g (Proteolytic act	

MSDS

Protease S "Amano"
(*Bacillus stearothermophilus*)

Powder

8.0

70°C

 $\geq 10,000$ u/g
(Proteolytic act)**MSDS**

Proleather FG
(*Bacillus subtilis*)

Powder

10.0

60°C

 $\geq 10,000$ u/g
(Proteolytic act)**MSDS**

Acylase

Acylase "Amano"
(*Aspergillus sp.*)

Powder

8.5

50°C

 $\geq 30,000$ u/g
(pH 8.0)**Catalogue****MSDS**

D-aminoacylase "Amano" *
(*E. coli*)

Powder lyophilized

8.0

40°C

 $\geq 5,000,000$ u/
(pH 8.0)**Catalogue****MSDS**

Other

Glucose Dehydrogenase
(*Bacillus sp.*)

Powder lyophilized

8.0

50°C

 $\geq 30,000$ u/g
(pH 8.0)**MSDS**

*under development

Each country has its own regulations on enzymes.

Please confirm fitness and regulations for your application to your products of our enzymes at your responsibility.



COPYRIGHT© 2003 AMANO ENZYME INC. ALL RIGHTS RESERVED.



CONTENTS

• About
Enzymes

• Product
Information

• Enzyme
Applications

• Quality
Assurance

• Enzyme
Wave

Company Information

Product Information

Product
I form

Select

Food Processing Use

For inquiries or sample request

Category	Products	Form	Opt. pH	Opt. Temp	Specification	Applic
Amylase	AMT 1.2L (<i>Microbacterium imperiale</i>) Catalogue MSDS	Liquid	6.5	50-55°C	S-Amylase ≥ 1,200 u/ml (pH 6.0)	Maltotriose Rich
	Transglucosidase L "Amano" (<i>Aspergillus niger</i>) Catalogue MSDS	Liquid	5.0	60°C	Transglucosidase ≥ 300,000 u/ml (pH 5.0)	Isomalto- oligosac Syrup
	Cyclodextrin glucanotransferase "Amano" (<i>Bacillus macerans</i>) Catalogue MSDS	Liquid	6.0	60-65°C	CGT-ase ≥ 600 u/ml (pH 5.5)	Production of cyclodextrin
Protease	Acid Protease II (<i>Rhizopus niveus</i>) Catalogue MSDS	Powder	3.0	45°C	Protease ≥ 15,000 u/g (pH 3.0)	Protein Hydrolysis
	Acid Protease A (<i>Aspergillus niger</i>) Catalogue MSDS	Powder	2.5	55°C	Protease ≥ 35,000 u/g (pH 3.0)	Protein Hydrolysis
	Newlase F (<i>Rhizopus niveus</i>) Catalogue MSDS	Powder	3.0	45°C	Protease ≥ 7,000 u/g (pH 3.0)	Protein Hydrolysis
	Protease A "Amano" 2 (<i>Aspergillus oryzae</i>) Catalogue MSDS	Powder	7.0	50°C	Protease ≥ 20,000 u/g (pH 7.0)	Protein Hydrolysis
	Protease M "Amano" (<i>Aspergillus oryzae</i>) Catalogue MSDS	Powder	4.5	50°C	Protease ≥ 5,500 u/g (pH 3.0)	Protein Hydrolysis

	Protease N "Amano" (<i>Bacillus subtilis</i>)	Powder	7.0	55°C	Protease ≥ 150,000 u/g (pH 7.0)	Protein Hydrolys
	Catalogue MSDS					
	Protease NL "Amano" (<i>Bacillus subtilis</i>)	Liquid	7.0	55°C	Protease ≥ 20,000 u/ml (pH 7.0)	Protein Hydrolys
	Catalogue MSDS					
	Protease P "Amano" 6 (<i>Aspergillus melleus</i>)	Powder	8.0	45°C	Protease ≥ 60,000 u/g (pH 8.0)	Protein Hydrolys
	Catalogue MSDS					
	Protease S "Amano" (<i>Bacillus stearothermophilus</i>)	Powder	8.0	70°C	Protease ≥ 10,000 u/g (pH 7.0)	Protein Hydrolys
	Catalogue MSDS					
	Proleather FG-F (<i>Bacillus subtilis</i>)	Granule	10.0	60°C	Protease ≥ 10,000 u/g (pH 10.0)	Protein Hydrolys
	Catalogue MSDS					
Lipase	Umamizyme (<i>Aspergillus oryzae</i>)	Powder	8.0	45°C	Peptidase ≥ 70 u/g (pH 7.0)	Bitterless Protein
	Catalogue MSDS					
	Peptidase R (<i>Rhizopus oryzae</i>)	Powder	7.0	45°C	Peptidase ≥ 420 u/g (pH 7.0)	Bitterless Protein
	Catalogue MSDS					
	Lipase A "Amano" 12 (<i>Aspergillus niger</i>)	Powder	6.5	45°C	Lipase ≥ 120,000 u/g (pH 6.0)	Fats & Oils Proc
	Catalogue MSDS					
	[Lipase AY "Amano" 30 (<i>Candida rugosa</i>)	Powder	7.0	45°C	Lipase ≥ 30,000 u/g (pH 7.0)	Fats & Oils Proc
	Catalogue MSDS					
	[Lipase F-AP15 (<i>Rhizopus oryzae</i>)	Powder	7.0	40°C	Lipase ≥ 150,000 u/g (FIP, pH 7.0)	Fats & Oils Proc
	Catalogue MSDS					
	[Lipase G "Amano" 50 (<i>Penicillium camembertii</i>)	Powder	5.0	40°C	Lipase ≥ 50,000 u/g (pH 5.6)	Fats & Oils Proc
	Catalogue MSDS					
	Lipase M "Amano" 10 (<i>Mucor javanicus</i>)	Powder	7.0	40°C	Lipase ≥ 10,000 u/g (pH 7.0)	Fats & Oils Proc
	Catalogue MSDS					
	Lipase R "Amano" (<i>Penicillium roqueforti</i>)	Powder	7.0	30°C	Lipase ≥ 900 u/g (pH 7.0)	Fats & Oils Proc
	Catalogue MSDS					

	Newlase F (<i>Rhizopus niveus</i>) Catalogue MSDS	Powder	7.0	40°C	Lipase ≥ 30,000 u/g (pH 7.0)	Fats & Oils Proc
Yeast Extract	Enzyme RP-1 (<i>Penicillium citrinum</i>) Catalogue MSDS	Powder	5.0	70°C	Nuclease ≥ 13,000 u/mg (pH 4.8)	Production of 5'
	Deamizyme 50000 (<i>Aspergillus melleus</i>) Catalogue MSDS	Powder	5.6	50°C	Deaminase ≥ 50,000 u/mg (pH 5.6)	Production of Inc
	YL-NL "Amano" (<i>Bacillus subtilis</i>) Catalogue MSDS	Liquid	6.0	50°C	Yeast Lysis activity ≥ 350 u/ml (pH 7.0)	Lyse the yeast c
	Glutaminase Daiwa C100S (<i>Bacillus subtilis</i>) Catalogue MSDS	Powder	6.0-7.0	65°C	Glutaminase 110GTU/g±9% (pH 6.0)	Production of L-t
Others	Hyderase (<i>Aspergillus niger</i>) Catalogue MSDS	Powder	6.0-8.0	40°C	Glucose oxidase ≥ 15,000 u/g (pH 7.0)	Removal of Oxy Baking
	Hyderase HC (<i>Aspergillus niger</i>) Catalogue MSDS	Powder	6.0-8.0	40°C	Glucose oxidase ≥ 1,500 u/g (pH 7.0) (High Catalase Content)	Removal of Oxy Baking
	Hyderase L (<i>Aspergillus niger</i>) Catalogue MSDS	Liquid	6.0-8.0	40°C	Glucose oxidase ≥ 10,000 u/ml (pH 7.0)	Removal of Oxy
	Dextranase L "Amano" (<i>Chaetomium erraticum</i>) Catalogue MSDS	Liquid	5.5	60°C	Dextranase ≥ 30,000 u/ml (pH 5.8)	Sugar Refining
	Lactase F "Amano" (<i>Aspergillus niger</i>) Catalogue MSDS	Powder	4.5	60°C	Lactase ≥ 14,000 u/g (pH 4.5)	Lactose Hydroly:
	Lactase Y "Amano" L (<i>Kluyveromyces Lactis</i>) Catalogue MSDS	Liquid	6.5	40°C	Lactase ≥ 10,000 u/g (pH 6.5)	Lactose Hydroly:

Each country has its own regulations on enzymes.

Please confirm fitness and regulations for your application to your products of our enzymes at your responsibility.



COPYRIGHT(C) 2003 AMANO ENZYME INC. ALL RIGHTS RESERVED.

The Wellcome Trust
Sanger Institute

Sanger Home | Acedb | YourGenome | Ensembl | Trace Server | Library
Info | Databases | Blast | Genomics | Infrastructure | HGP | CGP | Projects | Software | Teams | SearchData Release Policy | Conditions of L

Burkholderia cenocepacia

Projects Home

The Sanger Institute has been funded by Beowulf Genomics to sequence the genome of *Burkholderia cenocepacia*, in collaboration with Dr. Eshwar Mahenthiralingham of the School of Biosciences, Cardiff University, Prof. John Govan of the Department of Medical Microbiology, University of Edinburgh, Prof. C.A. Hart of the Department of Medical Microbiology, Royal Liverpool Hospital, and Prof. Peter Vandamme of the Laboratorium voor Microbiologie, Universiteit Gent, Belgium.

B. cenocepacia

Overview

Artemis

Blast Search

Sequence FTP

Related Projects

B. pseudomallei

Microbes Home

Search

FTP Site

Sitemap

Burkholderia cenocepacia (formerly *Burkholderia cepacia*, and before that *Pseudomonas cepacia*) is a Gram negative bacterium which is ubiquitous in the environment and may cause a number of diseases in plants. Human infection can be caused by *B. cepacia*, especially in patients with cystic fibrosis and chronic granulomatous disease, and is often fatal. The strain to be sequenced, J2315, is a genomovar III strain of the ET12 lineage, and is the index strain for transmission of this strain among CF patients in the UK. Strain J2315 has been deposited as LMG 16656 in the Belgium Coordinated Collection of Microorganisms.

Note that *B. cenocepacia* J2315 has recently been renamed from *B. cepacia* J2315. Information about this change is published in Vandamme *et al.* (2003) Res Microbiol. 154 91-96

Sequencing is now complete, and the sequences are available for searching on our Blast Server, or for download from our FTP site.

The genome totals 8.056 Mb in three chromosomes of 3.870, 3.217 and 0.876 Mb, and a plasmid of 92.7 kb, with a G+C content of approximately 66.9%.

Annotation of the sequence is ongoing, and the full annotation will be released upon publication. Please note that, although the sequence is finished, and we believe it to be accurate, it is possible that errors and missassemblies may remain. The sequence should be considered as preliminary until final publication. We would ask users of the data to read our Data release policy and Guidelines on use of data in publications

This sequencing centre plans on publishing the completed and annotated sequences in a peer-reviewed journal as soon as possible. Permission of the principal investigator should be obtained before publishing analyses of the sequence/open reading frames/genes on a chromosome or genome scale.

Please address all sequencing enquiries to: parkhill@sanger.ac.uk or barrell@sanger.ac.uk

DSMZ - List of Microbial Species: *Rhizopus javanicus* (Filamentous Fungi)**DSMZ**

© by DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany

Name	<i>Rhizopus javanicus</i> Takeda, see also: <i>Rhizopus oryzae</i> Went & Prinsen Gerligs
Strains	listed under other name: 2192 see: <i>Rhizopus oryzae</i>

DSMZMicroorganisms

DSM 2192 - *Rhizopus oryzae* (Filamentous Fungi)

© by DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany

Name	<i>Rhizopus oryzae</i> Went & Prinsen Gerligs (Filamentous Fungi)
DSM No.	2192
=	CBS 391.34
Information	<- CBS <- R. Nakazawa (<i>R. javanicus</i> , type strain). (Medium 129, 30°C)
Medium	129
Supplied as	actively growing culture (on agar or in liquid medium, depending on the strain)
Price	EURO 38 (non-profit making institutions), EURO 54 (other institutions): <u>Normal price.</u>

DSMZ

Microorganisms

DSMZ - List of Microbial Species: *Rhizopus delemar* (Filamentous Fungi)

© by DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany

Name	<i>Rhizopus delemar</i>
Strains	listed under other name: 853

DSMZ

Microorganisms

DSM 853 - *Rhizopus oryzae* (Filamentous Fungi)

© by DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany

Name	<i>Rhizopus oryzae</i> Went & Prinsen Gerligs (Filamentous Fungi)
DSM No.	853
	= ATCC 4858, CBS 327.47
Information	<- K. Kieslich <- ATCC <- A.J. Kluyver (<i>R. delemar</i>). Produces pectinase. (Medium 129, 30°C)
Medium	129
Supplied as	actively growing culture (on agar or in liquid medium, depending on the strain)
Price	EURO 38 (non-profit making institutions), EURO 54 (other institutions): <u>Normal price.</u>

DSMZ

Microorganisms

UNLOCKING THE
POWER OF ENZYME

ABOUT US PRODUCTS & SOLUTIONS DISCOVER BIOTECH OUR SCIENCE BIOTIMES PRESS ROOM INVESTOR ZONE CUSTOMER C
Our Solutions | Industry Calendar | Enzyme Quality Management

What is your area of interest?

Biocatalysis

What do you want to use our product for?

Commercial Products

Select specific product

Product



Commercially available products

Novozymes is offering a series of commercially available products for biocatalysis 'off the shelf'. By working with these products, researchers developing new processes for synthesis of new molecules, e. g. new drugs, or developing new bio-based processes to replace existing traditional processes have the assurance that the enzymes are available in large quantities if they make a hit. This assurance is based on Novozymes' unrivalled enzyme production capabilities. We produce enzymes by the tonne rather than the gram - economically!

This works to your advantage. All too often we have seen researchers waste months of research and valuable work because the enzyme on which they based their research was only available in minute, experimental quantities. The enzyme could not be upscaled and made available in sufficient quantities - or only at a prohibitively high, show-stopping cost.

This will not be the case if you work with our commercial product range:

Product

Novozym® 388
Lipozyme® RM IM
Novozym 735
Novozym CALB L
Novozym 435
Novozym 525 F
Lipozyme TL 100 L
Lipozyme TL IM
Novozym 539 HP F
Esperase® HP F
Subtilisin-A
Esperase HP F
Cryst. Porcine Trypsin
Novo 4500 K

Enzyme type

Rhizomucor miehei lipase
Rhizomucor miehei lipase
Candida antarctica lipase A
Candida antarctica lipase B
Candida antarctica lipase B
Candida antarctica lipase B
Lipase
Lipase
Protease
Protease
Protease
Protease
Trypsin



Site map | He

VISIT OUR CUSTOMERS

VISIT OUR CUSTOMERS

VISIT OUR CUSTOMERS

VISIT OUR CUSTOMERS

VISIT OUR CUSTOMERS

VISIT OUR CUSTOMERS



VISIT OUR CUSTOMERS

Detailed information on these products as well as further informative and supporting literature is available at the Customer Centre at Novozymes.com. If you do not already have access to the Customer Centre, please fill in the application form or contact us.

You can also place orders online through the Buyer's Centre section of the Customer Centre.

[Send to a friend »](#)

[Print version »](#)

© 2000-02 Novozymes A/S

Novozymes is a biotech-based world leader in enzymes and microorganisms.

[privacy](#)

UNLOCKING THE
FUTURE OF BIOTECH

ABOUT US PRODUCTS & SOLUTIONS DISCOVER BIOTECH OUR SCIENCE BIOTIMES PRESS ROOM INVESTOR ZONE CUSTOMER C
Our Solutions | Industry Calendar | Enzyme Quality Management

What is your area of interest?

Biocatalysis

What do you want to use our product for?

Commercial Products

Select specific product

Novozym 435



Novozym® 435

Novozym 435 is a lipase (lipase B) from *Candida antarctica* produced by submerged fermentation of a genetically modified *Aspergillus* microorganism and adsorbed on a macroporous resin.

Novozym 435 is probably the most commonly used lipase in biocatalysis. It is a highly versatile catalyst with activity towards a great variety of different substrates. The immobilized enzyme is a very thermostable and robust catalyst with activity in a number of organic solvents. It can be used both in batch and column reaction operations but is particularly well suited for use in fixed-bed reactors. Even though Novozym 435 is used in industrial processes as a catalyst in the synthesis of simple esters (including polyesters) and amides and in the regioselective synthesis of carbohydrate monoesters, it is primarily used as a highly enantioselective catalyst in the synthesis of optically active alcohols, amines and carboxylic acids.

[Send to a friend »](#)[Print version »](#)[Site map | He](#)[FOCUS](#)[SOLUTIONS](#)[DISCOVER](#)[BIOTIMES](#)[PRESS ROOM](#)[INVESTOR](#)

VISIT OUR CUSTOMERS

UNLOCKING THE
FUTURE OF BIOTECH

ABOUT US PRODUCTS & SOLUTIONS DISCOVER BIOTECH OUR SCIENCE BIOTIMES PRESS ROOM INVESTOR ZONE CUSTOMER C
Our Solutions | Industry Calendar | Enzyme Quality Management

What is your area of interest?

Biocatalysis

What do you want to use our product for?

Commercial Products


Select specific product

Novozym 525 F



Novozym® 525 F

Novozym 525 F is a lipase (lipase B) from *Candida antarctica* produced by submerged fermentation of a genetically modified *Aspergillus oryzae* microorganism. Novozym 525 F is a highly versatile catalyst with activity towards a great variety of different substrates. The enzyme is used in particular as a powerful enantioselective catalyst in the synthesis of optically active alcohols, amines and carboxylic acids.

[Send to a friend »](#)[Print version »](#)[Site map](#) | [Home](#)

Chemo-Enzymatic Syntheses of Both Enantiomers of Neodictyoprolenol and Neodictyoprolene; Possible Biosynthetic Intermediates of Sex Pheromones in Brown Algae

Yuuko Yamamoto, Yoshihiko Akakabe, Kenji Matsui, Hiroshi Shimidzu and Tadahiko Kajiwara*

Department of Biological Chemistry, Faculty of Agriculture, Yamaguchi University, Yamaguchi 753-8515, Japan. Fax: +81-839-33-5849. E-mail: kajiwara@ayu.agr.yamaguchi-u.ac.jp

* Author for correspondence and reprint requests

Z. Naturforsch. **54c**, 1027–1032 (1999); received June 7/July 12, 1999
Algae, Enantioselective Synthesis, Neodictyoprolene, Pheromones

Neodictyoprolenol [(–)-(S)-(1,5Z,8Z)-undecatrien-3-ol], dictyoprolenol [(–)-(S)-(1,5Z)-undecadien-3-ol] and their acetates neodictyoprolene [(+)-(S)-3-acetoxy-(1,5Z,8Z)-undecatriene] and dictyoprolene [(+)-(S)-3-acetoxy-(1,5Z)-undecadiene], which are interesting as possible biosynthetic intermediates of the sex pheromones (dictyopterene **B**, **C'** and **D'**) of brown algae, were synthesized by chemo-enzymatic methods through optical resolution of racemic neodictyoprolenol and dictyoprolenol using two lipases; Amano PS (*Pseudomonas* sp.) and Novozym 435® (*Candida* sp.). A combination of acylation of the alcohols and hydrolysis of the acetates by Novozym 435® produced neodictyoprolenol, neodictyoprolene, dictyoprolenol and dictyoprolene with high optical purities over 99% enantiomeric excess (*e.e.*).

This synthetic methods will make it easier to search these compounds in marine algae and to study their biosynthesis.

Enzymatic synthesis of derivatives of vitamin A in organic media

Thierry Maugard ^{*}, Marie Dominique Legoy

Laboratoire de Génie Protéique et Cellulaire, UPRES 2001, UFR Sciences et Technologie, Université de La Rochelle, Avenue Marillac,
17042 La Rochelle Cedex 1, France

Received 12 April 1999; accepted 18 June 1999

Abstract

The present article provides an enzymatic method of retinol esterification to reduce photodestruction and irritation problems characteristic of retinol. More specifically, it relates to a method of synthesising retinyl adipate, retinyl succinate, retinyl oleate and retinyl lactate greatly appreciated by cosmetic manufacturer. Desired compounds can be synthesised directly using *Candida antarctica* lipase and *Rhizomucor miehei* lipase in organic media. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: Retinol; Enzymatic synthesis; Lipase; Cosmetics

1. Introduction

Vitamin A (retinol) and derivatives (retinyl esters, retinoic acid, retinaldehyde), are of great commercial potential in cosmetics and pharmaceuticals such as skin care products. This water-insoluble vitamin has been shown to enhance gap junction communication in a dose-dependent manner, and may promote normal cell growth [1,2]. Vitamin A is necessary for maintenance of healthy epithelial tissue and can prevent the inception or progress of skin cancers by stimulating normal cell differentiation. Deficiency of vitamin A is usually accompanied by a number of severe diseases [3,4]. Consequently, in certain conditions extra amounts are needed in order to cure or prevent certain of

these diseases. But vitamin A is very unstable (readily oxidised in air and inactivated by UV light) and difficult to formulate satisfactorily for topical application. Moreover, vitamin A is irritating to the skin. Consequently, to reduce photodestruction and irritation problems, vitamin A may be stabilised by conversion to vitamin A ester (retinyl ester).

Various chemical methods have been published describing the synthesis of retinyl esters. However, retinol degradation often is important, leading to low yields. Moreover, these methods often have the disadvantage that they produce amounts of undesired residual chemical reactants in the final products such as salts.

In view of the attractiveness of retinyl esters as cosmetic agents, as therapeutic and nutritional additives, enzymatic methods can be suggested to produce retinyl esters by a “biological” process. Hydrolytic enzymes such as

^{*} Corresponding author. Tel.: +33-5-46-45-82-77; fax: +33-5-46-45-82-47; E-mail: tmaugard@univ-lr.fr

lipases in particular, have received special attention because of their effectiveness in regioselective and enantioselective esterifications and transesterifications of organic acids and alcohols in organic solvents [5,6]. Applications of such enzymatic reactions have been considered, for example, in the synthesis of flavor [7–9] and fatty-acid esters [10,11], in the regioselective acylation of carbohydrates [12], in the chemoselective acylation of amino-sugar derivatives [13,14] and in the synthesis of lactones [15], peptides [16], and chiral drugs [17].

The present article presents a preliminary study of the enzymatic retinol esterification using lipases in organic media.

2. Experimental

2.1. Biological and chemical material

Novozym® SP 435 (lipase from *Candida antarctica* immobilised on an acrylic resin), Lipozyme® (lipase from *Rhizomucor miehei* immobilised on an anionic macroporous resin, Duolite 568N), were from Novo Industries (Denmark). Porcine pancreatic lipase, *C. cylindracea* lipase and *Rhizopus arrhizus* lipase, were from Sigma (USA).

The solvents, all analytical grade, retinol, dimethyl adipate and dimethyl succinate, monomethyl succinate, succinic acid, methyl oleate and L-methyl lactate were from Sigma.

2.2. General procedure for the enzymatic reaction

Reactions were conducted in screw-cap glass vials containing 0.5 mmol of retinol **1**, 0.5 mmol of acyl donor, 100 mg of lipase and 0.25 g of 4-Å molecular sieves with 5 ml of solvent. The reaction mixture was stirred under positive argon pressure at 55°C and protected from light. After 50 h the biocatalyst and the molecular sieve were removed by filtration.

2.3. HPLC analysis

Analyses were performed with an HPLC system from Hewlett Packard (processor, pump, UV detector and injector model 1100, differential refractometer (Waters) model 410), equipped with an Ultrasep C18 (250 × 4 mm, 6 µ) reverse phase column from ICS, France. 25 µl of the appropriate dilution of the reaction mixture were injected. A mixture of methanol/water/acetic acid, 90/10/0.3 (v/v/v) was used as eluent at 40°C and a flow rate of 1 ml/min. Products were detected using a UV detector at 280 nm and a differential refractometer. The samples were quantitated by means of calibration curves with pure reagent.

2.4. Purification of reaction products

At the end of the reaction, the biocatalyst was removed by filtration and the solvent evaporated under reduced pressure. The remaining oil was purified by HPLC preparative using an Ultrasep C18 (250 × 8 mm, 6 µ) reverse phase column from ICS, France. A mixture of methanol/water/acetic acid, 90/10/0.3 (v/v/v) was used as eluent at 40°C and a flow rate of 3 ml/min. Products were detected using a UV detector at 280 nm.

2.5. Structural analysis

¹³C and ¹H Nuclear Magnetic Resonance spectra were recorded on a JEOL-JNM LA400 (400 MHz) spectrometer (Laboratoire Commun d'Analyse, Université de La Rochelle), with an internal reference of tetramethylsilane. IR spectra were recorded on a Perkin Elmer Paragon 1000PC instrument in NaCl plate. Mass spectra were recorded on a Varian MAT311 in the "Centre Régional de Mesure Physiques de L'Ouest" (CRMPO), Université de Rennes, France.

(Cad: carbon of adipate; Csu: carbon of succinate; Col: carbon of oleate; Cla: carbon of lactate.)

(Had: proton of adipate; Hsu: proton of succinate; Hol: proton of oleate; Hla: proton of lactate.)

2.5.1. Retinyl methyl succinate (3a)

IR: $\nu(\text{CH}) = 2800\text{--}2900\text{ cm}^{-1}$, $\nu(\text{=CH}) = 3100\text{ cm}^{-1}$, and $\nu(\text{CO--O ester}) = 1740\text{ cm}^{-1}$; ^{13}C NMR/DMSO, 6D (δ in ppm): 173 and 172.5 (2Co_{su} ester), 138.7, 137.3, 135.9 and 128.9 (C=), 137.2, 136.0, 130.3, 126.3, 125.8 and 125.14 (CH=), 60.5 (CH₂O), 39.5, 32.5 and 18.7 (CH₂ cyclohexenyl), 21.5 (CH₃), 28.8 (2 \times CH₃), 12.5 (2 \times CH₃ cyclohexenyl), 33.9 (C cyclohexenyl), 51.2 (CH₃O), 33.0, 33.3 (CH₂–CO_{su}), 23.8 (2 \times CH₂su); ^1H NMR/DMSO, 6D (δ in ppm): 0.98 (6H, s, 2 \times CH₃ cyclohexenyl), 1.42 (2H, m, CH₂ cyclohexenyl), 1.56 (2H, m, CH₂ cyclohexenyl), 1.99 (2H, m, CH₂ cyclohexenyl), 1.67 (3H, s, CH₃–C=), 1.84 (3H, s, CH₃–C=), 1.89 (3H, s, CH₃–C=), 4.68 (2H, d, CH₂O), 5.41 (1H, t, CH=), 5.62 (1H, t, CH=), 6.13 (2H, 2d, 2 \times CH=), 6.32 (1H, d, CH=), 6.66 (1H, dd, CH=), 2.3 (4H, m, 2 CH₂CO_{su}), 3.56 (3H, s, OCH₃su); C₂₅H₃₆O₄ requires M, 400.2613, found M⁺, 400.2618; m/z 400 (M⁺, 3.16%), 268 (M⁺–C₅H₈O₄, 4.17%), 43 (C₃H₇, 100%).

2.5.2. Retinyl succinate (3b)

IR: $\nu(\text{CH}) = 2800\text{--}2900\text{ cm}^{-1}$, $\nu(\text{=CH}) = 3100\text{ cm}^{-1}$, $\nu(\text{CO--O acid}) = 1725\text{ cm}^{-1}$ and $\nu(\text{CO--O ester}) = 1740\text{ cm}^{-1}$; ^{13}C NMR/DMSO, 6D (δ in ppm): 173 (CO_{su} ester), 170 (Co_{su} acid), 138.7, 137.3, 135.9 and 128.9 (C=), 137.2, 136.0, 130.3, 126.3, 125.8 and 125.14 (CH=), 60.5 (CH₂O), 39.5, 32.5 and 18.7 (CH₂ cyclohexenyl), 21.5 (CH₃), 28.8 (2 \times CH₃), 12.5 (2 \times CH₃ cyclohexenyl), 33.9 (C cyclohexenyl), 33.0, 33.3 (CH₂–CO_{su}), 23.8 (2 \times CH₂su); ^1H NMR/DMSO, 6D (δ in ppm): 0.98 (6H, s, 2 \times CH₃ cyclohexenyl), 1.42 (2H, m, CH₂ cyclohexenyl), 1.56 (2H, m, CH₂ cyclohexenyl), 1.99 (2H, m, CH₂ cyclohexenyl), 1.67 (3H, s, CH₃–C=), 1.84 (3H, s, CH₃–C=), 1.89 (3H, s, CH₃–C=), 4.68 (2H, d, CH₂O), 5.41 (1H, t, CH=), 5.62 (1H, t, CH=), 6.13


(2H, 2d, 2 \times CH=), 6.32 (1H, d, CH=), 6.66 (1H, dd, CH=), 2.3 (4H, m, 2 CH₂CO_{su}); C₂₄H₃₄O₄ requires M, 386.2456, found M⁺, 386.2462; m/z 386 (M⁺, 6.32%), 268 (M⁺–C₄H₆O₄, 5.22%), 43 (C₃H₇, 100%).

2.5.3. Retinyl methyl adipate (3c)

IR: $\nu(\text{CH}) = 2800\text{--}2900\text{ cm}^{-1}$, $\nu(\text{=CH}) = 3100\text{ cm}^{-1}$ and $\nu(\text{CO--O ester}) = 1737\text{ cm}^{-1}$; ^{13}C NMR/DMSO, 6D (δ in ppm): 173 and 172.5 (2CO_{ad} ester), 138.4, 137.3, 135.9 and 128.9 (C=), 137.2, 136.0, 130.3, 126.3, 125.8 and 125.14 (CH=), 60.5 (CH₂O), 39.5, 32.5 and 18.7 (CH₂ cyclohexenyl), 21.5 (CH₃), 28.8 (2 \times CH₃), 12.5 (2 \times CH₃ cyclohexenyl), 33.9 (C cyclohexenyl), 51.2 (CH₃O), 33.0, 33.3 (CH₂–CO_{ad}), 23.8 (2 \times CH₂ad); ^1H NMR/DMSO, 6D (δ in ppm): 0.98 (6H, s, 2 \times CH₃ cyclohexenyl), 1.42 (2H, m, CH₂ cyclohexenyl), 1.56 (2H, m, CH₂ cyclohexenyl), 1.99 (2H, m, CH₂ cyclohexenyl), 1.67 (3H, s, CH₃–C=), 1.84 (3H, s, CH₃ to C=), 1.89 (3H, s, CH₃–C=), 4.68 (2H, d, CH₂O), 5.41 (1H, t, CH=), 5.62 (1H, t, CH=), 6.13 (2H, 2d, 2 \times CH=), 6.32 (1H, d, CH=), 6.66 (1H, dd, CH=), 1.50 (4H, m, CH₂–CH₂ad), 2.3 (4H, m, 2 CH₂CO_{ad}), 3.56 (3H, s, OCH₃ad); C₂₇H₄₀O₄ requires M, 428.2926, found M⁺, 428.2935; m/z 428 (M⁺, 7.52%), 268 (M⁺–C₇H₁₂O₄, 3.26%), 43 (C₃H₇, 100%).

2.5.4. Retinyl L-lactate (3d)

IR: $\nu(\text{CH}) = 2800\text{--}2900\text{ cm}^{-1}$, $\nu(\text{=CH}) = 3100\text{ cm}^{-1}$, $\nu(\text{OH}) = 3400\text{ cm}^{-1}$ and $\nu(\text{CO--O ester}) = 1740\text{ cm}^{-1}$; ^{13}C NMR/DMSO, 6D (δ in ppm): 174.5 (CO_{la} ester), 138.7, 137.3, 135.9 and 128.9 (C=), 137.2, 136.0, 130.3, 126.3, 125.8 and 125.14 (CH=), 60.8 (CH₂O), 39.5, 32.5 and 18.7 (CH₂ cyclohexenyl), 21.5 (CH₃), 28.8 (2 \times CH₃), 12.5 (2 \times CH₃ cyclohexenyl), 33.9 (C cyclohexenyl), 65.9 (CH₃la), 20.4 (CH₃la); ^1H NMR/DMSO, 6D (δ in ppm): 0.98 (6H, s, 2 \times CH₃ cyclohexenyl), 1.42 (2H, m, CH₂ cyclohexenyl), 1.56 (2H, m, CH₂ cyclohexenyl), 1.99 (2H, m, CH₂ cyclohexenyl), 1.67 (3H, s, CH₃–C=), 1.84 (3H, s, CH₃–C=),



Scheme 1.

1.89 (3H, s, CH₃-C=), 4.71 (2H, d, CH₂O), 5.41 (1H, t, CH=), 5.62 (1H, t, CH=), 6.13 (2H, 2d, 2 × CH=), 6.32 (1H, d, CH=), 6.66 (1H, dd, CH=), 4.13 (1H, m, CH_{lact}), 1.23 (3H, d, CH₃_{lact}); C₂₃H₃₄O₃ requires M, 358.25078, found M⁺, 358.2522; *m/z* 358 (M⁺, 13.52%), 268 (M⁺-C₃H₆O₃, 52.87%), 45 (C₂H₅O₁, 100%), 43 (C₃H₇, 70.48%).

2.5.5. Retinyl oleate (3e)

IR: $\nu(\text{CH}) = 2800\text{--}2900\text{ cm}^{-1}$, $\nu(\text{=CH}) = 3100\text{ cm}^{-1}$ and $\nu(\text{CO-O ester}) = 1740\text{ cm}^{-1}$; ¹³C NMR/DMSO, 6D (δ in ppm): 174 (CO_{ol} ester), 138.7, 137.3, 135.9 and 128.9 (C=), 137.2, 136.0, 130.3, 126.3, 125.8 and 125.14 (CH=), 60.5 (CH₂O), 39.5, 32.5 and 18.7 (CH₂ cyclohexenyl), 21.5 (CH₃), 28.8 (2 × CH₃), 12.5 (2 × CH₃ cyclohexenyl), 33.9 (C cyclohexenyl), 130 (2 CH=CH_{ol}), 22.7–35.8 (14 CH₂_{ol}), 14 (CH₃_{ol}); C₃₈H₆₂O₂ requires M, 550.4749, found M⁺, 550.4754; *m/z* 550 (M⁺,

10.86%), 268 (M⁺-C₁₈H₃₄O₂, 4.17%), 43 (C₃H₇, 100%).

3. Results and discussion

The synthesis of retinyl methyl succinate **3a** from retinol **1** and dimethyl succinate **2a**, was chosen as the model reaction (Scheme 1). The *C. antarctica* lipase (Novozym[®]) and *R. miehei* lipase (Lipozyme[®]) were selected as catalysts because these immobilised enzymes are very efficient to the synthesis by reverse hydrolysis in organic media. The reaction was run in screw-cap glass vials containing 0.5 mmol of retinol **1**, 0.5 mmol of dimethyl succinate **2a**, 100 mg of lipase and 0.25 g of 4-Å molecular sieves with 5 ml of different solvents for 50 h at 55°C. From HPLC analysis, the decrease of retinol **1** concentration was seen to be concomitant with the synthesis of **3a**. In the absence of the enzyme, product **3a** did not appear. The

Table 1

Analytical yields of retinyl methyl succinate synthesis in different solvents

All the reactions were carried out at 55°C, with 100 mM of retinol, 100 mM of dimethyl succinate and 10 g/l of lipase in the presence of 4-Å molecular sieves (0.25 g/5 ml solvent). The degree of conversion was determined by HPLC on the base of the substrates' disappearance. No acylation was observed without lipase.

Solvent	Yield (%) with <i>C. antarctica</i> lipase	Yield (%) with <i>R. miehei</i> lipase	Reaction time (h)
Hexane	75	77	50
4-Methyl-2-pentanone	26	20	50
<i>t</i> -Amyl-alcohol	24	17	50
Dioxane	8	2	50
2-Butanone	0	0	50
Cyclohexanone	0	0	50

Table 2

Esterification of retinol with various acyl donors, catalysed by *R. miehei* lipase

All the reactions were carried out at 55°C, in hexane or in *t*-amyl-alcohol/hexane* (2/3, v/v), with 100 mM of retinol, 100 mM of acyl donor and 10 g/l of lipase in the presence of 4-Å molecular sieves (0.25 g/5 ml solvent). The degree of conversion was determined by HPLC on the base of the substrates' disappearance. No acylation was observed without lipase.

Acylating agent	Solvent	Yield after 50 h (%)	Product
Succinic acid	<i>t</i> -Amyl-alcohol/hexane*	4	Retinyl succinate 3b
Monomethyl succinate	<i>t</i> -Amyl-alcohol/hexane*	44	Retinyl methyl succinate 3a
Dimethyl adipate	Hexane	83	Retinyl methyl adipate 3c
Methyl lactate	Hexane	86	Retinyl L-lactate 3d
Methyl oleate	Hexane	90	Retinyl oleate 3e

isolation, purification and characterisation of product **3a** confirmed that **3a** corresponded to retinyl methyl succinate.

The different yields obtained depending on the nature of solvents and on the nature of lipase, are reported in Table 1. When cyclohexanone and 2-butanone are used as reaction solvents, lipase activity is not detectable. In all the other tests, retinol esterification occurs with variable yields, depending on the solvent and lipase used. The highest yield is obtained in hexane. Three other commercial lipases (porcine pancreatic lipase, *C. cylindracea* lipase and *Rhi. arrhizus* lipase), were tested under the same conditions but no significant reaction was observed.

In addition, using *R. miehei*, other acyl donors were used to produce a wide variety of retinyl esters of great interest in cosmetic formulations. The results are presented in Table 2. Reactions with succinic acid and monomethyl succinate are carried out in *t*-amyl-alcohol/hexane system (2/3, v/v), in order to solubilise acyl donors which are insoluble in hexane. With the exception of monomethyl succinate, we observed excellent yields (more than 80%) for all the methyl ester donors. On the other hand, with succinic acid and monomethyl succinate, we found a lower reactivity than with the methyl ester donors.

In order to improve the system performance and to shift the reaction equilibrium towards synthesis, all the reactions were run in *t*-amyl-alcohol at 90°C under reduced pressure. The

co-produced methanol was then removed more rapidly than under atmospheric pressure. We used *C. antarctica* lipase. This one has already shown its efficiency in similar conditions. However, in these conditions, the retinol degradation increases more rapidly than the retinol acylation. Indeed, retinol methyl succinate formation is competitive with the destruction of retinol characterised by the production of several compounds more hydrophilic than retinol.

4. Conclusion

An efficient method has been developed for the lipase-catalysed synthesis of retinyl esters such as retinyl succinate, retinyl adipate, retinyl oleate and retinyl lactate. To our knowledge, this enzymatic process is the first that describes retinyl esters synthesis with such excellent yields. In addition, the retinyl lactate produced may be considered as an excellent carrier of lactic acid. Indeed, in the presence of esterase-type epidermal enzymes, they can undergo hydrolysis, thus releasing lactic acid progressively. Derived from fruit and dairy products, alpha hydroxy acids are widely included in cosmetics as exfoliants. The most commonly used are lactic acid and glycolic acid, both of which seem to exert slight but significant effects in reducing skin discolorations and roughness when applied in a cream [18,19]. Significant irritation is often associated with the use of alpha hydroxy acids alone. Esters of retinol and alpha

hydroxy acids are unusually effective as skin conditioners, with significant reductions in the irritation problems characteristic of retinol and alpha hydroxy acids in nonesterified form [20].

Acknowledgements

The generous gift of *C. antarctica* lipase and *R. miehei* lipase by Novo-Nordisk is gratefully acknowledged.

References

- [1] J.S. Bertram, Cancer prevention by retinoids and carotenoids: proposed role of gap junctional communication, in: M.M. Jacobs (Ed.), Vitamins and Minerals in the Prevention and Treatment of Cancer, CRC Press, Boca Raton, 1991, pp. 31–50.
- [2] M.D. Karen Laszlo Keller, A. Neil, M.D. Fenske, J. Am. Acad. Dermatol. 39 (1998) 611–625.
- [3] W.H. Sebrell, R.S. Harris (Eds.), The Vitamins, 2nd edn., Vol. 1 (descriptive information on the chemistry physico-chemical properties and physiology of Vitamin A), Academic Press, NY, 1967.
- [4] R.J. Sokol, Annu. Rev. Nutr. 8 (1988) 351–373.
- [5] J.S. Dordick, Principles and applications of nonaqueous enzymology, in: H.W. Blanch, D.S. Clark (Eds.), Applied Biocatalysis, Vol. 1, Marcel Dekker, New York, 1991, pp. 1–52.
- [6] A. Zaks, A.M. Klibanov, Proc. Natl. Acad. Sci. U.S.A. 82 (1985) 3192–3196.
- [7] C. Marlot, G. Langrand, C. Triantaphylides, J. Baratti, Biotechnol. Lett. 1 (1985) 211–216.
- [8] A. Manjon, J.L. Iborra, A. Arocas, Biotechnol. Lett. 13 (1991) 339–344.
- [9] G. Carta, J.L. Gainer, M.E. Gibson, Enzyme Microb. Technol. 14 (1992) 904–910.
- [10] S.-W. Cho, J.S. Rhee, Biotechnol. Bioeng. 41 (1993) 204–210.
- [11] A. Zaidi, J.L. Gainer, G. Carta, Biotechnol. Bioeng. 48 (1995) 601–605.
- [12] S. Riva, F. Secundo, Chimicaoggi 6 (1990) 9–16.
- [13] T. Maugard, M. Remaud-Simeon, D. Petre, P. Monsan, Tetrahedron 53 (1997) 7587–7594.
- [14] T. Maugard, M. Remaud-Simeon, D. Petre, P. Monsan, Tetrahedron 53 (1997) 5185–5194.
- [15] C.-S. Chen, S.M. Wu, G. Girdaukas, C.J. Sih, J. Am. Chem. Soc. 109 (1987) 2812–2817.
- [16] A.L. Margolin, A.M. Klibanov, J. Am. Chem. Soc. 109 (1987) 3802–3804.
- [17] A.L. Margolin, Enzyme Microb. Technol. 15 (1993) 266–280.
- [18] E.C. Stern, Urol. Cutaneous Rev. 50 (1946) 106–107.
- [19] E.J. Scott Van, R.J. Yu, Arch. Dermatol. 110 (1974) 586–590.
- [20] T.J., Bigger, S. Nacht, S. Ng, International Patent Publication WO/97/20812, 1996.


UNLOCKING THE
MAGIC OF NATURE

ABOUT US | PRODUCTS & SOLUTIONS | DISCOVER BIOTECH | OUR SCIENCE | BIOTIMES | PRESS ROOM | INVESTOR ZONE | CUSTOMER C
Facts | Sales Offices | Visit our Sites | **Publications** | Novozymes TV | Management | Corporate Governance | Corporate Responsibility



Demerger document

- » The Novozymes Report
- » Annual Reports
- » Environmental & Social Reports
- » Site Reports
- » The Zymes - Shareholders Magazine
- » Enzymes at work
- » Why do we need enzymes?
- » Nature makes the enzymes
- We put them to work
- » Unlocking the magic of nature
- » **Demerger document**



In February 1999, Novo Nordisk announced its plans to demerge the company's activities within health care and enzymes for industrial use into separately listed legal entities.

The decision was taken at an extraordinary general meeting on 13 November 2000 in Copenhagen, Denmark. The demerger took effect retroactively as of 1 January 2000.

Download PDF version of Demerger Document

[Send to a friend »](#)

[Print version »](#)

© 2000-02 Novozymes A/S. Novozymes is a biotech-based world leader in enzymes and microorganisms.

privacy

Site map | He

Site map | He

Site map | He

Site map | He

Site map | He

Site map | He

Site map | He

The Novo Group
commitment to
Development



Aktieudbuddet
tegnet mere e

Novozymes A/
the Copenhagen
Exchange

Site map | He

Site map | He





PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bc

Search PubMed for [] Go Clear
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

1: J Biol Chem. 1989 Dec 15;264(35):21146-52.

Related Articles, Links

FREE full text article at
www.jbc.org

Characterization of the relA1 mutation and a comparison of relA1 with new relA null alleles in Escherichia coli.

Metzger S, Schreiber G, Aizenman E, Cashel M, Glaser G.

Department of Cellular Biochemistry, Hebrew University Hadassah Medical School, Jerusalem, Israel.

The most widely studied "relaxed" mutant of the relA locus, the relA1 allele, is shown here to consist of an IS2 insertion between the 85th and 86th codons of the otherwise wild-type relA structural gene, which normally encodes a 743-amino acid (84 kDa) protein. The RelA protein is a ribosome-dependent ATP:GTP (GDP) pyrophosphoryltransferase that is activated during the stringent response to amino acid starvation and thereby occasions the accumulation of guanosine 3',5'-bispyrophosphate (ppGpp). We propose that the IS2 insertion functionally splits the RelA protein into two (alpha and beta) peptide fragments which can complement each other in trans to yield residual ppGpp synthetic activity; neither fragment shows this activity when expressed alone. Cell strains with a single copy relA null allele show physiological behavior that is much the same as relA1 mutant strains. Both relA1 and relA null strains accumulate ppGpp during glucose starvation and do not accumulate ppGpp during the stringent response. The presence of ppGpp in verifiable relA null strains is interpreted as unequivocal evidence for an alternate route of ppGpp synthesis that exists in addition to the relA-dependent reaction.

PMID: 2556396 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Linkage Map of *Escherichia coli* K-12, Edition 10: The Traditional Map

MARY K. B. BERLYN*

Department of Biology and School of Forestry and Environmental Studies, Yale University,
 New Haven, Connecticut 06520-8104

INTRODUCTION	814
MAP UNITS	821
NOMENCLATURE.....	821
Gene Symbol Convention.....	821
The Issue of Stability	821
ACKNOWLEDGMENTS	888
REFERENCES	889

INTRODUCTION

Previously, Berlyn et al. (323) presented the traditional map, the EcoMap physical map, and a map by Singer and Low showing the distribution of the Gross-Singer transposon set around the chromosome. The map in this paper is a revision of that traditional map of *Escherichia coli* K-12, the linkage map of known genes and other functional sites (Fig. 1), and the physical map, EcoMap 10, of Kenneth Rudd is presented in the companion article (3763a).

The linkage map in this presentation includes genes located primarily by restriction, sequence, and cotransduction data reported in the literature and databases. It uses coordinates based on the complete sequence released by the Blattner laboratory. Obviously, the sequence is now the major resource for placing genes on the map. In some regions the placement represents a shift from the edition 9 map, which was based on coordinates of Rudd's EcoMap 7 composite of sequenced genes and regions (27, 33, 395, 568, 569, 926, 3308, 3465, 4127, 4128), placed on the physical map of *Escherichia coli* (2291, 3763b) by restriction and sequence comparisons. Those map positions were based on the results in the literature and on EcoMap and GenBank database entries. EcoMap 10 coordinates are of course also based on the completed sequence, and cross-consulting this summary map and the EcoMap that follows should be straightforward.

The linkage map of Fig. 1 includes 2,220 genes and about 40 other chromosomal markers, such as phage attachment sites, defective-phage elements, replication origins and termini, and other features traditionally included on the published linkage map. It does not include open reading frames (ORFs) lacking evidence for expression, with unknown functions or putative functions inferred by sequence homologies only. A few exceptions occur for *Salmonella* genes where the inference is strong that they are also expressed in *E. coli*. The ORFs not included

in this map can be found on EcoMap 10. The Fig. 1 map places the genes that can be found in sequence annotation and EcoMap 10 on the right side of the line. On the left side are genes not present on physical maps or the sequence, and in most cases these are not connected to a specific point on the axis to indicate that the localization is only approximate. As in previous editions of the *E. coli* linkage map (187, 188, 189, 190, 190a, 323, 4368, 4369, 4370, 4371), an asterisk indicates that the gene is not precisely located with respect to near neighbors and parentheses indicate that the location is even more uncertain and that the gene is located only within that general region. I have been very conservative about removing these from the map; even though the usefulness of some of these may be quite limited, there will probably be cases where the old, sometimes poorly characterized phenotype may be helpful in ascribing functions and phenotypic effects to ORFs. Also shown on the left side in boldface followed by colons are operon names that are distinct from any gene name within the operon and termination and attachment sites. The arrows indicate the direction of transcription and span genes included within a transcription unit.

Updates of map information are available in electronic form from several sites. These include the *E. coli* Genetic Stock Center's (CGSC's) World Wide Web server at URL <http://cgsc.biology.yale.edu>, which provides an interface for querying the database and retrieving formatted reports about genes, map regions, strains, and mutations, etc. (323a); the National Center for Biotechnology Information ftp site for EcoSeq and EcoMap, ncbi.nlm.nih.gov/repository/Eco/EcoMap7; the Colibri map at <http://www.pasteur.fr/Bio/Colibri.html>, the ECDC map at <http://susi.bio.uni-giessen.de/ecdc.html>, the site for the sequencing project at the University of Wisconsin, <http://www.genetics.wisc.edu>, a gene-protein database, <http://www.mbl.edu/html/ecoli.html>, Genome Information Broker at <http://mol.genes.nig.ac.jp/ecoli>, and others. See also Rudd (3673). The references attempt to document map information, the basic definition of the gene's function, and expression information and do not include information relating to detailed physical structure, active site in vitro mutagenesis, or enzyme

* Mailing address: 355 Osborn Memorial Laboratories, 165 Prospect St., Box 208104, Yale University, New Haven, CT 06520-8104. Phone: (203) 432-9997. Fax: (203) 432-3854. E-mail: mary.berlyn@yale.edu.

FIG. 1. Linear drawing of circular linkage map of *E. coli* K-12. Symbols are defined in Table 1. Arrows show the direction of transcription. Where T-bars are used to display groups of genes, the length of the T shows the approximate length and position of the group in terms of the map coordinates, allowing visual ordering of closely packed groups.






FIG. 1—Continued.








FIG. 1—Continued.



mechanism. Earlier map papers contain additional references for some of the loci (188–190a, 323).

MAP UNITS

Since the 1976 recalibration of the linkage map in terms of minutes required for time of entry of markers in interrupted conjugation experiments, the standard representation of the map has used the basic units of minutes and a total length of 100 minutes (190a). This has been a convenient and accepted coordinate system for the map, and although the current map units are based on restriction and sequence data rather than time of entry, we retain the term minute for 1/100 of the length of the chromosome. Both the CGSC database and EcoMap use as "left endpoints" the counterclockwise boundary of the coding region, and genes in Fig. 1 are placed approximately at these coordinates, with the higher-resolution map of Rudd (3763a) providing more exact placement, showing nucleotide and minute coordinates for the physically mapped genes.

NOMENCLATURE

Gene Symbol Convention

The standard genetic nomenclature for *E. coli* is that of Demerec et al. (1016), as subsequently amended through use, and as described in Instructions to Authors for the *Journal of Bacteriology* (see also reference 3821). This map, like those preceding it, follows those nomenclatural conventions. Accordingly, we have adhered to a three-letter lowercase mnemonic symbol, with an uppercase letter added when there are two or more genes in that mnemonic category. If authors have added an uppercase letter for a gene in a single-instance category, we have used that published four-letter symbol. For attachment sites and noncoding features of the chromosome, etc., the same standard has not been used, and we have continued to use the variable-length symbols historically applied to these sites. We have continued the convention proposed for sites of termination of replication and repetitive sequences, by using italicized symbols with the first letter uppercase.

The Issue of Stability

Many names have been changed by investigators since the 1990 map was published. When those changes were part of a systematic revision of nomenclature (often aimed at clarifying usage and resolving conflicts) for a group of related genes and were in compliance with the current *E. coli* gene nomenclature system, or were changed for compelling mnemonic reasons or for resolution of redundancy or conflict, also in conformance with the standard system, we have adopted those changes. We have not adopted and we wish to discourage changes of valid preexisting names proposed by authors simply because they

believe that theirs is a symbol signifying a more apt or accurate mnemonic. For example, a previously published name based on the pathway or phenotype is valid and should not, simply as a matter of course, be replaced by an alternate mnemonic based on the name of the enzyme that the gene codes for once that functional information has been determined. In general, the stability of a name has more value than improved nuances. In a few cases, we have been compelled to use a new name, despite the apparent validity of the original name, simply because the new name has been widely adopted in the literature. In a number of cases, a new gene has been assigned a symbol which has already been used or which is simultaneously proposed for another gene, with the two mnemonics having entirely different meanings. These names have had to be resolved, usually by changing the newer assignment. In a few cases, a uniquely named gene has been shown later to belong to a category for which a symbol already exists, and the latter symbol has been used instead of the earlier assignment. There is one case in this paper where use of a symbol already assigned to a different gene was strongly preferred by authors, and I was very reluctant to suggest a new symbol for the earlier, published gene name to the earlier authors, since that symbol has been used in a number of publications; for the interim I have broken convention to assign the newer genes temporary symbols with asterisks (*gsp**), in order to show them on the map and in the hope of resolving that naming with the usual precedence custom in the near future. Some synonymy is unavoidable, since a gene under study may be named and described in print before its identity to a known gene is discovered. However, a common practice in the recent literature seems to allow publication of an author's preliminary name for a gene even if its identity to a known gene has been discovered before publication, and that practice creates unnecessary synonymy. Alternate gene symbols are listed in Table 1, and Table 2 provides an alphabetized list of such symbols with cross-references to the symbols used in Table 1.

There is a standing tradition of coordinating gene symbols between the CGSC and the *Salmonella* Genetic Stock Center to avoid the assignment of the same symbol to different genes in the two organisms and the assignment of different symbols to homologous genes, insofar as this coordination is feasible. We have not, however, changed names of *E. coli* genes in order to extend this tradition to other bacteria or other organisms. The desirability and feasibility of uniform nomenclature conventions for all bacteria or other microbial groupings are currently only topics of discussion and conjecture, and changes to enhance similarities in an ad hoc, piecemeal fashion seem counterproductive at the present time. Readers are reminded that symbol changes create discontinuity with previous literature concerning genes, with even more serious ramifications for allele designations, since unique allele numbers are assigned on the basis of the three letter mnemonic and changes in a symbol may necessitate renumbering of alleles as well.

TABLE 1. *E. coli* genes and replication- or phage-related sites^a

Gene symbol	Map location (min)	Mnemonic for symbol	Synonyms and gene product—enzyme, RNA, or phenotype affected	CGSC no. ^d	References ^c
<i>aarF</i>	86.6	Aminoglycoside acetyltransferase regulator	<i>yigQR</i> ; regulator of 2'-N-acetyltransferase; involved in respiratory cofactor ubiquinone production	53879	2696
<i>aas</i>	64.1	Acyl-ACP synthase	2-Acyl-glycerophosphoethanolamine acyltransferase; acyl-ACP synthetase; salvage pathway for reacylation; inner membrane; bifunctional for turnover/incorporation	29780	1831, 1972
<i>aat</i>	20.0	Amino acyl-tRNA-protein transferase	Aminoacyl-tRNA-protein-transferase (EC 2.3.2.6)	1054	4045
<i>abpS</i>	63.5	Arg binding protein	Low-affinity transport system for arginine and ornithine; periplasmic binding protein	18562	664
<i>abs</i>	94.1	Antibiotic sensitivity	Sensitivity and permeability to antibiotics and dyes	18559	763
<i>accA</i>	4.5	Acetyl-CoA carboxylase	Acetyl-CoA carboxylase α -carboxyltransferase subunit; (EC 6.4.1.2)	29829	2536, 2537
<i>accB</i>	73.4	Acetyl-CoA carboxylase	<i>fabE</i> ; acetyl-CoA carboxylase, biotin carboxyl carrier protein (EC 6.4.1.2)	796	2537, 2712, 3057, 4302, 4616
<i>accC</i>	73.4	Acetyl-CoA carboxylase	<i>fabG</i> ; acetyl-CoA carboxylase, biotin carboxylase (BC) subunit (EC 6.4.1.2)	29834	2315, 2537, 3253
<i>accD</i>	52.4	Acetyl-CoA carboxylase	<i>dedB</i> , <i>usg</i> ; acetyl-CoA carboxylase β -carboxyltransferase subunit (EC 6.4.1.2)	28570	2534, 2537, 3081
<i>acd</i>	65.1	Acetaldehyde-CoA deHase	Acetaldehyde-CoA dehydrogenase (EC 1.2.1.10)	1053	764
<i>aceA</i>	90.8	Acetate	<i>icl</i> ; isocitrate lyase (EC 4.1.3.1); acetate utilization	1052	2744, 2811, 3674, 754, 841
<i>aceB</i>	90.8	Acetate	<i>mas</i> ; malate synthase A (EC 4.1.3.2)	1051	2744, 591, 592, 840, 841
<i>aceE</i>	2.7	Acetate	<i>aceE1</i> ; pyruvate dehydrogenase (decarboxylase component) E1p; (EC 1.2.4.1); acetate requirement	1050	1553, 1555, 1556, 1558, 2934, 4150, 4204, 655
<i>aceF</i>	2.7	Acetate	<i>aceE2</i> ; pyruvate dehydrogenase (dihydrolipoyltransacetylase component) E2p (EC 1.6.4.3, EC 2.3.1.12); acetate requirement	26530	1554, 2934, 655
<i>aceK</i>	90.9	Acetate	Isocitrate dehydrogenase kinase/phosphatase	17770	1343, 1897, 2274, 2418, 2419, 753, 842, 754
<i>ackA</i>	52.0	Acetate kinase	Acetate kinase (EC 2.7.2.1); mutants fluoroacetate resistant	1048	1548, 2094, 2512, 2812, 530, 224
<i>ackB</i>	39.9	Acetate kinase	Acetate kinase activity* (EC 2.7.2.1)	1047	3379
<i>acnA</i>	28.8	Aconitase	Aconitase A (EC 4.2.1.2)	28218	3542, 3543
<i>acnB</i>	2.8	Aconitase	Aconitase B (EC 4.2.1.2)	36955	1538, 2141, 474
<i>acpD</i>	31.9	Acyl carrier protein	ACP phosphodiesterase	52896	1256, 394
<i>acpP</i>	24.8	Acyl carrier protein	ACP (acyl carrier protein)	31871	2183, 3621
<i>acpS</i>	58.2	Acyl carrier protein	<i>dpi</i> ; holo-ACP synthase (EC 2.7.8.7)	32953	2397, 2400, 2401, 3471, 4340
<i>acpX</i>	43.9	Acyl carrier protein	<i>acpS</i> ; originally thought to be holo-ACP synthase; perhaps cryptic second gene or regulator	1046	3471
<i>acrA</i>	10.4	Acridine	<i>Mb</i> , <i>lir</i> , <i>mbl</i> , <i>mtcA</i> , <i>sipB</i> ; AcrAB efflux system effects Mar multiple resistance	1045	1291, 1726, 2684, 2685, 3282, 3563, 4723, 808
<i>acrB</i>	10.4	Acridine	<i>acrE</i> ; AcrAB system has major role in Mar multiple resistance to NAL, TET, AMP, etc.; beware renamings of <i>acrE</i> , <i>acrB</i> , and <i>envC</i>	35806	1291, 2684, 2685, 3282, 4723
<i>acrC</i>	4.5	Acridine	Sensitivity to acriflavine; transmembrane protein	1044	3104
<i>acrD</i>	55.7	Acridine	Sensitivity to acriflavine	35697	3188
<i>acrE</i>	73.5	Acridine	<i>envC</i> ; anomolous cell division; chain formation; splits cross-wall to form new poles; see <i>acrB</i>	813	2133, 2255, 2256, 2516, 3702
<i>acrF</i>	73.6	Acridine	<i>envD</i> ; encodes lipoprotein with signal peptide; osmotically remedial envelope defect	33608	2255, 2256, 2516, 2684
<i>acrR</i>	10.5	Acridine	Regulatory protein for <i>acrA</i> and <i>acrB</i>	35809	2686
<i>acs</i>	92.3	Acetyl-CoA synthetase	Acetyl CoA synthetase 2 (EC 6.2.1.1)	34317	2367, 395
<i>ada</i>	49.7	Adaptive (response)	O ⁶ -methylguanine-DNA methyltransferase, inducible; DNA repair against methylating and alkylating agents; transcription factor	1043	1184, 2001, 2158, 2410, 2489, 2698, 2760, 2967, 3078, 3094, 3795, 3812, 3815, 3957, 3958, 3959, 4330, 4385
<i>add</i>	36.6	Adenine deaminase	Adenosine deaminase (EC 3.5.4.4); mutants affect growth on deoxyadenosine in <i>purA</i> , <i>B</i> mutants	1042	2024, 765

Continued on following page

TABLE 1—Continued

Gene symbol	Map location (min)	Mnemonic for symbol	Synonyms and gene product—enzyme, RNA, or phenotype affected	CGSC no. ^d	References ^e
<i>dinG</i>	17.9	Damage inducible	LexA regulated (SOS) repair enzyme	31247	2318, 2518, 2519
<i>dinI</i>	24.1	Damage inducible	Multicopy suppresses phenotype of cold-sensitive <i>dinD</i> filamentous mutation	53428	4877
<i>dinY</i>	41.9	Damage inducible	Repair gene	36880	3412
<i>dipZ</i>	94.0	Disulfide isomerase	<i>cycZ</i> , <i>dsbD</i> , <i>cutA2</i> ; may be involved in cytochrome maturation, see <i>ccm</i> genes; affects disulfide binding	34213	1921, 268, 2910, 874, 875
<i>djlA</i>	1.2	DnaJ-like	Proposed to dock and interact with variety of membrane proteins; mutants rapidly accumulate suppressors	51192	2188, 769, 770
<i>dksA</i>	3.5	<i>dnaK</i> suppressor	<i>msmA</i> ; high copy suppresses <i>muk</i> and TS growth and filamentation of <i>dnaK</i> mutant	30521	2122, 4842
<i>dld</i>	47.9	D-Lactate dehydrogenase	<i>ldh</i> ; D-lactate dehydrogenase (EC 1.1.1.28); vinylglycolate resistance, FAD enzyme	852	2058, 3768, 4012, 626
<i>dmsA</i>	20.3	DMSO reductase	DMSO reductase subunit A, anaerobic	17710	353, 354, 3749
<i>dmsB</i>	20.3	DMSO reductase	DMSO reductase subunit B; apparent Fe-S binding domain; anaerobic	31724	354, 3749
<i>dmsC</i>	20.3	DMSO reductase	DMSO reductase subunit C, membrane bound	31727	354, 3749
<i>dnaA</i>	83.6	DNA	DNA biosynthesis; initiation; binding protein	851	1372, 1630, 1631, 1872, 1873, 1875, 2160, 2235, 2272, 2275, 2474, 2927, 2985, 2986, 3051, 3261, 3400, 3541, 3569, 3570, 3809, 3813, 4087, 4300, 4301, 4611, 471, 4910, 1790
<i>dnaB</i>	91.9	DNA	<i>groP</i> , <i>grpA</i> , <i>grpD</i> ; DNA biosynthesis; chain elongation	850	1406, 2259, 2545, 3126, 3333, 3804, 3951, 4449, 561
<i>dnaC</i>	99.1	DNA	<i>dnaD</i> ; DNA biosynthesis; initiation and chain elongation	849	2786, 3125, 3753
<i>dnaE</i>	4.4	DNA	<i>polC</i> , <i>sdgC</i> (suppressor of <i>dnaG</i> mutation); DNA polymerase III, α -subunit	373	1239, 1240, 2194, 296, 3889, 4016, 4424, 4708, 4761
<i>dnaG</i>	69.2	DNA	<i>dnaP</i> , <i>parB</i> , <i>sdgA</i> ; primase; primer synthesis for leading- and lagging-strand synthesis	847	1531, 2671, 2672, 2673, 3055, 3113, 3125, 3216, 3753, 4105, 4376, 4449, 4578, 4766, 579
<i>dnaI</i>	40.3	DNA	DNA biosynthesis	846	334a
<i>dnaJ</i>	0.3	DNA	<i>groP</i> , <i>grpC</i> ; chain elongation; stress-related DNA biosynthesis, responsive to heat shock; chaperone with DnaK	845	1406, 2069, 231, 3256, 3257, 3258, 3333, 3804, 4252, 4293, 4830, 972
<i>dnaK</i>	0.3	DNA	<i>gro</i> , <i>groP</i> , <i>groPAB</i> , <i>groPC</i> , <i>groPF</i> , <i>grpC</i> , <i>grpF</i> , <i>seg</i> ; stress-related heat-shock DNA biosynthesis, ATP-regulated binding and release of polypeptide substrates; HSP-70-type molecular chaperone, with DnaJ	844	1143, 1186, 1187, 1875, 2120, 227, 2837, 3035, 3158, 3257, 3332, 3807, 4252, 567, 3697
<i>dnaL</i>	28.9	DNA	DNA biosynthesis	843	3982
<i>dnaN</i>	83.6	DNA	DNA biosynthesis; sliding clamp subunit, required for high processivity; DNA polymerase III β subunit	842	132, 1630, 2194, 2235, 3261, 3626, 3627, 3809, 3810, 3813, 4910, 564
<i>dnaQ</i>	5.1	DNA	<i>mutD</i> ; DNA polymerase III ϵ -subunit; streptomycin, azaserine resistant; 3' to 5' proofreading, <i>lexA</i> regulon	840	1047, 1113, 1239, 1706, 1804, 2194, 2730, 2783, 3209, 3568, 3903, 4052, 779, 855, 856, 4331
<i>dnaT</i>	99.1	DNA	Primasomal protein i	839	2786, 2787, 3125
<i>dnaX</i>	10.6	DNA	<i>mutH</i> , <i>dnaZ</i> ; subunit of DNA polymerase III holoenzyme; DNA elongation factor III; τ and γ subunits	838	1264, 1265, 1266, 1669, 1845, 2194, 2228, 2229, 2283, 2421, 2471, 2472, 2511, 2731, 3043, 3753, 398, 399, 4485, 4486, 4885, 701, 912, 913
<i>dppA</i>	79.8	Dipeptide permease	<i>alu</i> , <i>tpp?</i> ; uptake of dipeptides	35111	3294, 4, 4576
<i>dppB</i>	79.8	Dipeptide permease	Uptake of dipeptides	33771	4, 4118, 4576
<i>dppC</i>	79.8	Dipeptide permease	Uptake of dipeptides	33768	4, 4118, 4576
<i>dppD</i>	79.8	Dipeptide permease	Uptake of dipeptides	33765	4, 4118, 4576
<i>dppF</i>	79.7	Dipeptide permease	Uptake of dipeptides	33752	4

Continued on following page

TABLE 1—Continued

Gene symbol	Map location (min)	Mnemonic for symbol	Synonyms and gene product—enzyme, RNA, or phenotype affected	CGSC no. ^d	References ^c
<i>murA</i>	71.8	Murein	<i>mrhA</i> , <i>murZ</i> ; UDP- <i>N</i> -acetylglucosamine enolpyruvyl transferase (EC 2.5.1.7); phosphomycin resistance	33518	2773, 3548, 4571, 527
<i>murB</i>	89.9	Murein	UDP- <i>N</i> -acetylglucosaminyl-3-enolpyruvate reductase (EC 1.1.1.158)	34141	308, 3548, 395
<i>murC</i>	2.2	Murein	L-Alanine adding enzyme	476	1183, 1893, 2021, 2071, 2675, 2676
<i>murD</i>	2.1	Murein	UDP- <i>N</i> -acetylmuramoyl-L-alanine:D-glutamate ligase (EC 6.3.2.9)	30450	1894, 2889, 2890, 3533
<i>murE</i>	2.0	Murein	<i>meso</i> -Diaminopimelate adding enzyme	475	1183, 2071, 2889, 2923, 4349
<i>murF</i>	2.0	Murein	<i>mra</i> ; D-alanyl:D-alanine adding enzyme	474	101, 1183, 2071, 2889, 3368
<i>murG</i>	2.1	Murein	UDP-NAc-glucosamine: NAc-muramyl-(pentapeptide) pyrophosphoryl-undecaprenol NAc-glucosamine transferase	473	1893, 2071, 2889, 2895, 3820
<i>murH</i>	99.3	Murein	Terminal stage in peptidoglycan synthesis, incorporating disaccharide peptide units into wall	18136	3204, 900
<i>murI</i>	89.7	Murein	<i>mbrC</i> , <i>dga</i> , <i>glr</i> ; glutamate racemase (EC 5.1.1.3); D-glutamate synthesis, essential for peptidoglycan	29401	1076, 1077, 1078, 212, 2584, 4900, 523
<i>mutG</i>	43.7	Mutator	Mutation causes high C-to-T mutation in second C of CCAGG; near but distinct from <i>usr</i> ; provisionally termed <i>mutG</i>	28933	3766
<i>mutH</i>	64.0	Mutator	<i>mutR</i> , <i>prv</i> ; methyl-directed mismatch repair; see also <i>dnaX</i>	471	1222, 1484, 1485, 1706, 4052, 4488, 856
<i>mutL</i>	94.7	Mutator	Methyl-directed mismatch repair	470	1088, 1222, 139, 1706, 4052, 4487, 4489, 4490, 856
<i>mutM</i>	82.1	Mutator	<i>fpg</i> ; repair; GC to TA; formamidopyrimidine-DNA glycosylase	18133	1221, 1489, 1706, 2434, 3717, 4052, 4128, 416, 417, 568, 599, 805, 856
<i>mutS</i>	61.5	Mutator	<i>ant?</i> <i>fdv</i> (formate dehydrogenase 2?); methyl-directed mismatch repair	469	1222, 1706, 2587, 3615, 3853, 4052, 4488, 4802, 856
<i>mutT</i>	2.4	Mutator	AT to GC transversions	468	1288, 1706, 2729, 338, 39, 4052, 856
<i>mutY</i>	66.8	Mutator	<i>micA</i> ; GC to TA transversions; adenine glycosylase, G-A repair	18130	1706, 2922, 3172, 3576, 4052, 856, 4476, 4477
<i>nac</i>	44.4	Nitrogen assimilation control	Regulatory gene, binding <i>ntnC</i>	50182	3073
<i>nadA</i>	16.8	Nicotinamide adenine dinucleotide	Quinolinate synthetase, A protein	467	1257, 2819, 4246
<i>nadB</i>	58.4	Nicotinamide adenine dinucleotide	Quinolinate synthetase, B protein	466	1257, 3015, 4379, 3963
<i>nadC</i>	2.5	Nicotinamide adenine dinucleotide	<i>nic</i> ; quinolinate phosphoribosyl transferase	465	1553, 1556, 1558, 2415, 2416, 3689
<i>nadE</i>	39.2	Nicotinamide adenine dinucleotide	<i>efg</i> , <i>ntnL</i> ; NAD synthetase, ammonia dependent	28576	4752, 70
<i>nagA</i>	15.1	<i>N</i> -Acetylglucosamine	<i>N</i> -acetylglucosamine-6-phosphate deacetylase (EC 3.5.1.25)	464	2491, 3402, 3454–3456, 4599, 4726
<i>nagB</i>	15.1	<i>N</i> -Acetylglucosamine	<i>glmD</i> ; glucosamine-6-phosphate deaminase (EC 5.3.1.10)	463	3402, 3454, 3455, 3456, 3464, 3709, 4601, 4726
<i>nagC</i>	15.1	<i>N</i> -Acetylglucosamine	<i>nagR</i> ; transcriptional regulator of <i>nag</i> operon	31479	3402, 3455, 3456, 3457, 4593, 3458
<i>nagD</i>	15.1	<i>N</i> -Acetylglucosamine	Function unknown, expressed as part of <i>nag</i> operon	36240	3455
<i>nagE</i>	15.2	<i>N</i> -Acetylglucosamine	<i>pstN</i> ; <i>N</i> -acetylglucosamine-specific enzyme II of phosphotransferase system	462	2048, 2491, 3401, 3402, 3454, 3709, 4599, 4726, 4600
<i>nalB</i>	60.2	Nalidixic acid	Sensitivity to nalidixic acid (NAL)	460	1626
<i>nalD</i>	89.2	Nalidixic acid	NAL sensitivity; NAL and glycerol penetration	18124	1827
<i>nanA</i>	72.6	<i>N</i> -Acetylneuraminate	<i>N</i> -Acetylneuraminate lyase (aldolase) (EC 4.1.3.3)	17647	2782, 3270, 4585
<i>nanT</i>	72.6	<i>N</i> -Acetylneuraminate	Sialic acid transport	18121	2782, 4585
<i>napA</i>	49.5	NitrAte reductase, periplasmic	Nitrate reductase homolog	36550	1537, 739, 740
<i>napB</i>	49.5	NitrAte reductase, periplasmic	Cytochrome <i>c</i> homolog	36560	1537, 739, 740
<i>napC</i>	49.5	NitrAte reductase, periplasmic	Cytochrome <i>c</i> homolog	36566	1537, 739, 740
<i>napD</i>	49.6	NitrAte reductase, periplasmic	Unknown function, <i>nap</i> operon	36547	1537, 739, 740

Continued on following page

TABLE 1—Continued

Gene symbol	Map location (min)	Mnemonic for symbol	Synonyms and gene product—enzyme, RNA, or phenotype affected	CGSC no. ^a	References ^c
<i>rcsA</i>	43.6	Regulation capsule synthesis	Positive regulatory gene for capsule (colanic acid) synthesis; two regulatory proteins from the same gene	17980	1045, 1471, 4243, 4244, 4441
<i>rcsB</i>	49.9	Regulation capsule synthesis	Positive regulatory gene for capsule (colanic acid) synthesis; when overexpressed, restores <i>ftsZ84</i> growth on low-salt medium	17977	1413, 1471, 2188, 4099, 4243–4245, 504
<i>rcsC</i>	49.9	Regulation capsule synthesis	Negative regulatory gene for capsule (colanic acid) synthesis, controls sliminess; contains TerE; probable histidine kinase	17974	1471, 2188, 3037, 4099, 4245, 504
<i>rcsF</i>	4.7	Regulation capsule synthesis	Overexpression confers mucoid phenotype, increases capsule synthesis; restores colony formation of <i>ftsZ84</i> mutants on low salt	29845	1412
<i>rdgA</i>	16.1	RecA-dependent growth	Dependence of growth upon <i>recA</i> gene product	17971	1317
<i>rdgB</i>	67.0	RecA-dependent growth	Dependence of growth and viability upon <i>recA</i>	17968	784
<i>recA</i>	60.8	Recombination	<i>srf</i> , <i>lexB</i> , <i>umuD</i> , <i>zab</i> ; general recombination and DNA repair; pairing and strand exchange; role in cleavage of LexA repressor, SOS mutagenesis	312	1104, 1123, 1239, 1654, 1803, 2177, 2574, 2575, 3833, 3991, 4395, 4604, 4749, 495, 497, 64, 671, 759
<i>recB</i>	63.6	Recombination	<i>rorA</i> ; recombination and repair; RecBCD enzyme (exonuclease V) subunit	311	1107, 1180, 1249, 1743, 3870, 4365, 690, 759, 4785a, 406a
<i>recC</i>	63.7	Recombination	Recombination and repair; RecBCD enzyme (exonuclease V) subunit	310	1107, 1252, 1262, 1743, 3870, 4365, 690, 759, 4785a
<i>recD</i>	63.6	Recombination	<i>hopE</i> ; recombination and repair; RecBCD enzyme (exonuclease V) α -subunit	4975	1248, 3187, 348, 4365, 759, 87
<i>recE</i>	30.5	Recombination	Recombination and repair; in prophage <i>rac</i> locus; degrades one strand 5'–3' in duplex DNA; exonuclease VIII	309	1181, 1287, 2086, 2087, 362, 4751, 748, 759, 760
<i>recF</i>	83.6	Recombination	<i>uvrF</i> ; recombination and repair	308	10, 11, 132, 1517, 152, 2709, 3626, 3627, 379, 3846, 3847, 3848, 4611, 671, 759
<i>recG</i>	82.4	Recombination	<i>spoV?</i> branch migration of Holliday junctions, junction-specific DNA helicase (see <i>nuvABC</i>)	307	1281, 2100, 2593, 2596, 2597, 2598, 2599, 3086, 4721, 59
<i>recJ</i>	65.4	Recombination	Single-stranded DNA-specific exonuclease, 5'–3'	17965	2638, 2639, 2640, 2641, 2642, 4511, 759
<i>recN</i>	59.3	Recombination	<i>radB</i> ; <i>lexA</i> regulon; recombination and repair	10872	2600, 2924, 3425, 3748, 3861, 3862
<i>recO</i>	58.2	Recombination	Conjugational recombination and repair; DNA-binding protein; RecA-like strand assimilation	17962	2297, 2657
<i>recQ</i>	86.3	Recombination	Conjugational recombination and repair, presynaptic stage of recombination; <i>lexA</i> regulon; RecQ helicase	17959	1923, 2883, 3123, 3124, 58, 926
<i>recR</i>	10.6	Recombination	Recombination and DNA repair	31049	2723
<i>recT</i>	30.4	Recombination	Locus in defective prophage <i>rac</i> ; activated by <i>sbcA</i> mutation; DNA-annealing protein	32070	1612, 2298, 2378, 761
<i>relA</i>	62.7	Relaxed	Required for ppGpp synthesis during stringent response to amino acid starvation; ATP:GTP 3'-pyrophosphotransferase (EC 2.7.6.5)	306	1237, 1441, 1442, 2913, 2914, 2915, 3070, 3606, 3934, 4310, 4807a, 924
<i>relB</i>	35.4	Relaxed	Stringent/relaxed response; regulation of RNA synthesis	305	1042, 266
<i>relE</i>	35.4	Relaxed	Function unknown	17956	1411, 266
<i>relF</i>	35.4	Relaxed	Function unknown; overproduction lethal	17953	1411, 266
<i>relX</i>	62.8	Relaxed	Control of ppGpp synthesis	304	3346
<i>rep</i>	85.3	Replicase	<i>dasC</i> , <i>mbrA</i> , <i>mmrA</i> ; Rep helicase, a single-stranded DNA-dependent ATPase	303	1086, 1428, 257, 2613, 342, 343, 4388, 58, 610, 926
<i>rer</i>	89.9	Resistance to radiation	Resistance to UV and gamma radiation	302	4179
<i>rfaB</i>	81.9	Rough	<i>waaB</i> ; UDP-galactose: (glucosyl)lipopolysaccharide-1,6-galactosyltransferase	17617	3361, 3522, 3726, 3921, 4128, 867
<i>rfaC</i>	81.8	Rough	<i>waaC</i> ; LPS core biosynthesis; proximal hexose; UDP-galactose: (glucosyl)LPS-1,6-galactosyltransferase	300	280, 3362, 3726, 3920, 3921, 4128, 702

Continued on following page

TABLE 2—Continued

Synonym	Gene symbol from Table 1	Synonym	Gene symbol from Table 1
<i>lig</i>	<i>ligA</i>	<i>msmC</i>	<i>cspE</i>
<i>lipP</i>	<i>nike?</i>	<i>msp</i>	<i>arcA</i>
<i>lir</i>	<i>acrA</i>	<i>msaA</i>	<i>cmk</i>
<i>livR</i>	<i>lrp</i>	<i>msbB</i>	<i>deaD</i>
<i>lky</i>	<i>tolA, B</i>	<i>msuA</i>	<i>dadX?</i>
<i>lkyA</i>	<i>tolB</i>	<i>msyA</i>	<i>hns</i>
<i>lnt</i>	<i>cutE</i>	<i>mtcA</i>	<i>acrA</i>
<i>lolB</i>	<i>hemM</i>	<i>mtcB</i>	<i>tolC</i>
<i>LopC</i>	<i>clpX</i>	<i>muc</i>	<i>lon</i>
<i>LopP</i>	<i>clpP</i>	<i>mukA</i>	<i>tolC</i>
<i>lov</i>	<i>argS</i>	<i>mukE</i>	<i>kicA</i>
<i>lovB</i>	<i>alaS</i>	<i>mukF</i>	<i>kicB</i>
<i>loxB</i>	<i>attP1, P7</i>	<i>murZ</i>	<i>murA</i>
<i>lrs</i>	<i>lrp</i>	<i>mutA</i>	<i>glyV</i>
<i>lss</i>	<i>lrp</i>	<i>mutC</i>	<i>glyW</i>
<i>lstR</i>	<i>lrp</i>	<i>mutD</i>	<i>dnaQ</i>
<i>luxH-like</i>	<i>ribB</i>	<i>mutH</i>	<i>dnaX</i>
<i>lysTα, β, γ</i>	<i>lysT, W, Y</i>	<i>mutR</i>	<i>topB</i>
<i>mac</i>	<i>maa</i>	<i>mutU</i>	<i>uvrD</i>
<i>malB</i>	<i>malK</i>	<i>nivA</i>	<i>fpr</i>
<i>maoA</i>	<i>tynA</i>	<i>nivC</i>	<i>emrE</i>
<i>maoB</i>	<i>feaR</i>	<i>nagR</i>	<i>nagC</i>
<i>mas</i>	<i>aceB</i>	<i>nalA</i>	<i>gyrA</i>
<i>mazE</i>	<i>chpR</i>	<i>nalC</i>	<i>gyrB</i>
<i>mazF</i>	<i>chpA</i>	<i>nalD</i>	<i>gyrB</i>
<i>Mb</i>	<i>acrA</i>	<i>nam</i>	<i>pncA</i>
<i>mhf</i>	<i>lrp</i>	<i>narA</i>	<i>moaA</i>
<i>mbl</i>	<i>acrA</i>	<i>narB</i>	<i>mobA</i>
<i>mbrA</i>	<i>rep</i>	<i>narC</i>	<i>narG</i>
<i>mcb</i>	<i>pmbA</i>	<i>narD</i>	<i>modC, E, F</i>
<i>mclA</i>	<i>rscA</i>	<i>narR</i>	<i>nurL, X</i>
<i>mdaA</i>	<i>nfsA</i>	<i>nbp</i>	<i>fis</i>
<i>mdfA</i>	<i>cmr</i>	<i>ncf</i>	<i>hemB</i>
<i>mdrA</i>	<i>cydC</i>	<i>neaA</i>	<i>rpsQ</i>
<i>mdrH</i>	<i>cydC</i>	<i>neck</i>	<i>rimK</i>
<i>mec</i>	<i>dcm</i>	<i>nfrC</i>	<i>rffE</i>
<i>mel-4</i>	<i>melB</i>	<i>nfsB</i>	<i>nfnB</i>
<i>mel-7</i>	<i>mclA</i>	<i>nfsI</i>	<i>nfnB</i>
<i>meoA</i>	<i>ompC</i>	<i>nfxA</i>	<i>gyrA</i>
<i>met-1</i>	<i>metB</i>	<i>nfxC</i>	<i>marA</i>
<i>metM</i>	<i>metL</i>	<i>nfxD</i>	<i>parE</i>
<i>metTB</i>	<i>metU</i>	<i>nic</i>	<i>nadB</i>
<i>metZβ</i>	<i>metV</i>	<i>nirA</i>	<i>fir</i>
<i>mglD</i>	<i>galS</i>	<i>nirR</i>	<i>fnr</i>
<i>mglP</i>	<i>mglA, C</i>	<i>nitA</i>	<i>rho</i>
<i>mhpS</i>	<i>mhpD</i>	<i>nitB</i>	<i>rpoB</i>
<i>micA</i>	<i>mutY</i>	<i>nlp</i>	<i>sfsB</i>
<i>mic</i>	<i>dgsA</i>	<i>nlpE</i>	<i>cutF</i>
<i>nlpA</i>	<i>lpp</i>	<i>nmpA</i>	<i>pst, pstS</i>
<i>nmrA</i>	<i>rep, rhlB?</i>	<i>nmpB</i>	<i>phoR</i>
<i>nni</i>	<i>manC</i>	<i>norA</i>	<i>gyrA</i>
<i>modR</i>	<i>modE</i>	<i>norB</i>	<i>marA?</i>
<i>molA</i>	<i>malM</i>	<i>nov</i>	<i>cls</i>
<i>momR</i>	<i>oxyR</i>	<i>ntr</i>	<i>nfnB</i>
<i>mon</i>	<i>mreB</i>	<i>ntrA</i>	<i>rpoN</i>
<i>mopA</i>	<i>groS</i>	<i>ntrB</i>	<i>glnL</i>
<i>mopB</i>	<i>groL</i>	<i>ntrC</i>	<i>glnG</i>
<i>mor</i>	<i>oxyR</i>	<i>ntrL</i>	<i>nadE</i>
<i>motD</i>	<i>flhN</i>	<i>nucR</i>	<i>deoR</i>
<i>mpt</i>	<i>manX, Z</i>	<i>nupA</i>	<i>tsx</i>
<i>mra</i>	<i>murF</i>	<i>nur</i>	<i>rpoS</i>
<i>mrba</i>	<i>murA</i>	<i>nusD</i>	<i>rho</i>
<i>mrc</i>	<i>lpcB</i>	<i>nusE</i>	<i>rpsJ</i>
<i>mrsA</i>	<i>glmM</i>	<i>oldB</i>	<i>fadB</i>
<i>mrsC</i>	<i>hflB</i>	<i>ole</i>	<i>fadR</i>
<i>msgA</i>	<i>ftsN</i>	<i>ompB</i>	<i>envZ, ompR</i>
<i>msmA</i>	<i>dksA</i>	<i>ompE</i>	<i>phoE</i>
<i>msmB</i>	<i>cspC</i>	<i>ompH</i>	<i>hlpA</i>

Continued on following page

TABLE 2—Continued

Synonym	Gene symbol from Table 1	Synonym	Gene symbol from Table 1
<i>tabD</i>	<i>rpoB</i> , C	<i>umpA</i>	<i>lgt</i>
<i>talA</i>	<i>alaT</i>	<i>umuA</i>	<i>lexA</i>
<i>talD</i>	<i>alaU</i>	<i>umuB</i>	<i>recA</i>
<i>tasC</i>	<i>aspT</i>	<i>uncA-I</i>	<i>atpA-I</i>
<i>tau</i>	<i>tus</i>	<i>ups?</i>	<i>prfA</i>
<i>tcp</i>	<i>rimJ</i>	<i>ura</i>	<i>car</i>
<i>tdc</i>	<i>tdcB</i>	<i>uraP</i>	<i>upp</i>
<i>tgs</i>	<i>crp</i> , <i>purU</i>	<i>usg</i>	<i>accD</i>
<i>tgtB</i>	<i>gltT</i>	<i>usgA</i>	<i>gntT</i>
<i>tgtC</i>	<i>gltU</i>	<i>uvm</i>	<i>umuC</i> , D
<i>tgtE</i>	<i>gltV</i>	<i>uvr502</i>	<i>uvrD</i>
<i>thdB</i>	<i>fudR</i>	<i>uvrE</i>	<i>uvrD</i>
<i>thdF</i>	<i>trmE</i>	<i>uvrF</i>	<i>recF</i>
<i>thiA</i>	<i>thiE</i> , F, G, H	<i>vacB</i>	<i>rrn</i>
<i>thiJ</i>	<i>thiL</i>	<i>val-act</i>	<i>valS</i>
<i>thiN</i>	<i>thiJ</i>	<i>valTα</i>	<i>valT</i>
<i>thrD</i>	<i>thrA</i>	<i>valTβ</i>	<i>valZ</i>
<i>thyR</i>	<i>deoB</i> , C	<i>valUα</i>	<i>valU</i>
<i>ildE</i>	<i>pmbA</i>	<i>valUγ</i>	<i>valY</i>
<i>tlhI</i>	<i>tlhA</i>	<i>virR</i>	<i>hns</i>
<i>tlr</i>	<i>deoB</i> , C	<i>visA</i>	<i>hemH</i>
<i>tls</i>	<i>aspS</i>	<i>visB</i>	<i>ubiH</i>
<i>tmuA</i>	<i>folA</i>	<i>vir</i>	<i>fabF</i>
<i>tnaC</i> , L	<i>tna</i> leader	<i>waaA</i>	<i>kdtA</i>
<i>tnaR</i>	<i>tnaA</i>	<i>waaB</i>	<i>rfaB</i>
<i>toc</i>	<i>tolC</i>	<i>waaC</i>	<i>rfaC</i>
<i>tol-2</i>	<i>tolA</i>	<i>waaE</i>	<i>rfaE</i>
<i>tol-3</i>	<i>tolB</i>	<i>waaF</i>	<i>rfaF</i>
<i>tolF</i>	<i>ompF</i>	<i>waaG</i>	<i>rfaG</i>
<i>tolG</i>	<i>ompA</i>	<i>waaI</i>	<i>rfaI</i>
<i>tolM</i>	<i>cmiB</i>	<i>waaJ</i>	<i>rfaJ</i>
<i>tolP</i>	<i>tolQ?</i>	<i>waaK</i>	<i>rfaK</i>
<i>tolZ</i>	<i>hflB</i>	<i>waaL</i>	<i>rfaL</i>
<i>tonA</i>	<i>fluA</i>	<i>waaM</i>	<i>htrB</i>
<i>topX</i>	<i>hns</i>	<i>waaN</i>	<i>mltA</i>
<i>tos</i>	<i>prfC</i>	<i>waaP</i>	<i>rfaP</i>
<i>tpo</i>	<i>envZ</i>	<i>waaQ</i>	<i>rfaQ</i>
<i>tpp</i>	<i>dppA?</i>	<i>waaS</i>	<i>rfaS</i>
<i>tpp-75</i>	<i>deoA</i>	<i>waaU</i>	<i>rfaK</i>
<i>tpx</i>	<i>ahpC</i>	<i>waaY</i>	<i>rfaY</i>
<i>tre</i>	<i>treA</i>	<i>waaZ</i>	<i>rfaZ</i>
<i>trkB</i>	<i>kefB</i>	<i>wbbH</i>	<i>rfe</i>
<i>trkC</i>	<i>kefC</i>	<i>wcaN</i>	<i>galF</i>
<i>trkE</i>	<i>sapD</i>	<i>wec</i>	<i>rff</i>
<i>trpP</i>	<i>tnaB</i>	<i>wecA</i>	<i>rfe</i>
<i>trpX</i>	<i>miaA</i>	<i>wecB</i>	<i>rffE</i>
<i>tryD</i>	<i>trpE</i>	<i>weeA</i>	<i>tolC</i>
<i>tryp-4</i>	<i>trpE</i>	<i>witA</i>	<i>kgiP</i>
<i>tsnC</i>	<i>trxA</i>	<i>wzxB</i>	<i>rfbX</i>
<i>tsp</i>	<i>prc</i>	<i>xerA</i>	<i>argR</i>
<i>tss</i>	<i>asnS</i>	<i>xerB</i>	<i>pepA</i>
<i>tsu</i>	<i>rho</i>	<i>xonA</i>	<i>sbcB</i>
<i>utr</i>	<i>fadL</i>	<i>xprA</i>	<i>dsbC</i>
<i>tut</i>	<i>ompA</i>	<i>xprB</i>	<i>xerD</i>
<i>tyrTα</i> , β	<i>tyrT</i> , V	<i>xylT</i>	<i>xylF</i>
<i>uar</i>	<i>prfA</i>	<i>zab</i>	<i>recA</i>
<i>uidP</i>	<i>uidB</i>	<i>zfaA</i>	<i>csrA</i>
<i>umg</i>	<i>ptsG</i>		

^a Temporary names (y names, *orfs*) are not included.

ACKNOWLEDGMENTS

This work was supported by the National Science Foundation grants BIR9315421 and BIR9010005.

I thank Stanley Letovsky and Peter Kalamarides for their work in implementation and system administration for the database, which allowed direct retrieval of all the various Table 1 data and the drawing of map segments directly from the CGSC database. Even so, extensive

layout work was required for the final figure, and I am indebted to Elise Low for her skill and patience in doing that during all the revisions that went into this version of the map. A number of people have worked tirelessly to get this paper together. Special thanks again go to Elise Low for editing graphics files and for unending proofreading and cross-checks on the tables and references in addition to the layout work. Completion of this task owes much to her skill and her dedication to accuracy and timeliness and to Peter Kalamarides for

writing and executing scripts that brought the tables together at every critical juncture. Special thanks also go to Linda Mattice and Narinder Whitehead for valuable help with proofreading and tracking down publications for both versions and for keeping the stock center on course during this work. I'm especially grateful to all of the above for their unstinting efforts at deadline-approaching time. I thank Graeme Berlyn for sharing and helping with use of his computer setup during the printing operation, and also James Bryan for table formatting help in the earlier edition. Brooks Low and Kenn Rudd coauthored the 1996 edition of the map and therefore were major contributors to this map as well; I thank them for that collaboration and their continued help and support. Kenn Rudd and I have attempted to keep this map and the physical map compatible in terms of names and annotations; Kenn has been crucial in keeping me updated with regard to his new information, displaying his characteristic dedication and generosity with regard to his sequence-to-gene detection and expertise. Special thanks this year go to Edward Adelberg for supplying the database with bibliographic updates based on his scanning and annotation of Medline references in the context of his own bibliographic database of current papers in *E. coli* genetics developed and kept up to date over the past 2 1/2 years. They have been enormously helpful in our attempts to keep the CGSC database current with respect to literature on *E. coli* genes. These maps continue the tradition of *E. coli* K-12 linkage map editions so capably constructed by Barbara J. Bachmann over the past 20 years and that of her predecessors for *E. coli* maps dating back to 1958. Our debt to these previous maps is obvious. The accuracy of the great majority of gene positions on the map is completely indebted to the American and Japanese sequencing projects, and for this particular version of the map, especially to GenBank access to the complete sequence submitted by Fred Blattner and his colleagues as part of the Genome Sequencing project in Wisconsin. And of course we are indebted to numerous scientists who have provided helpful map-related information in the form of discussion, corrections of database inaccuracies, preprints, and personal communications.

REFERENCES

- Aasland, R., J. Coleman, A. L. Holck, C. L. Smith, C. R. H. Raetz, and K. Kleppe. 1988. Identity of the 17-kilodalton protein, a DNA-binding protein from *Escherichia coli*, and the *firA* gene product. *J. Bacteriol.* 170: 5916-5918.
- Abailou, H., J. Pommier, S. Benoit, G. Giordano, and M.-A. Mandrand-Berthelot. 1995. Expression and characterization of the *Escherichia coli* *fdo* locus and a possible role for aerobic formate dehydrogenase. *J. Bacteriol.* 177:7141-7149.
- Abdulrashid, N., and D. P. Clark. 1987. Isolation and genetic analysis of mutations allowing the degradation of furans and thiophenes by *Escherichia coli*. *J. Bacteriol.* 169:1267-1271.
- Abouhamad, W. N., M. D. Manson, M. M. Gibson, and C. F. Higgins. 1991. Peptide transport and chemotaxis in *Escherichia coli* and *Salmonella typhimurium*: characterization of the dipeptide permease (Dpp) and the dipeptide-binding protein. *Mol. Microbiol.* 5:1035-1047.
- Abraham, J. M., C. S. Freitag, J. R. Clements, and B. I. Eisenstein. 1985. An invertible element of DNA controls phase variation of type 1 fimbriae of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 82:5724-5727.
- Abraham, S. N., J. D. Goguen, D. Sun, P. Klemm, and E. H. Beachey. 1987. Identification of two ancillary subunits of *Escherichia coli* type 1 fimbriae by using antibodies against synthetic oligopeptides of *fim* gene products. *J. Bacteriol.* 169:5530-5535.
- Abril, N., C. Hera, E. Alejandre, J. A. Rafferty, G. P. Margison, and C. Pueyo. 1994. Effect of *ogt* expression on mutation induction by methyl-, ethyl- and propylmethanesulphonate in *Escherichia coli* K12 strains. *Mol. Gen. Genet.* 242:744-748.
- Ackerman, R. S., N. R. Cozzarelli, and W. Epstein. 1974. Accumulation of toxic concentrations of methylglyoxal by wild-type *Escherichia coli* K-12. *J. Bacteriol.* 119:357-362.
- Adachi, H., T. Ohta, and H. Matsuzawa. 1987. A water-soluble form of penicillin-binding protein 2 of *Escherichia coli* constructed by site-directed mutagenesis. *FEBS Lett.* 226:150-154.
- Adachi, T., M. Mizuuchi, A. Robinson, E. Appella, M. H. O'Dea, M. Gellert, and K. Mizuuchi. 1987. DNA sequence of the *E. coli* *grb* gene: application of a new sequencing strategy. *Nucleic Acids Res.* 15:771-784.
- Adachi, T., R. Mizuuchi, R. Menzel, and M. Gellert. 1984. DNA sequence and transcription of the region upstream of the *E. coli* *grb* gene. *Nucleic Acids Res.* 12:6389-6395.
- Adam, E., M. R. Volkert, and M. Blot. 1998. Cytochrome c biogenesis is involved in the transposon Tn5-mediated bleomycin resistance and the associated fitness effect in *Escherichia coli*. *Mol. Microbiol.* 28:15-24.
- Adams, B. S., and R. B. Helling. 1970. Nalidixic acid-resistant auxotrophs of *Escherichia coli*. *J. Bacteriol.* 104:1027-1030.
- Adams, D. E., and S. C. West. 1996. Bypass of DNA heterologies during RuvAB-mediated three- and four-strand branch migration. *J. Mol. Biol.* 263:582-596.
- Adams, M. D., L. M. Wagner, T. J. Graddis, R. Landick, T. K. Antonucci, A. L. Gibson, and D. L. Oxender. 1990. Nucleotide sequence and genetic characterization reveal six essential genes for the LIV-I and LS transport systems of *Escherichia coli*. *J. Biol. Chem.* 265:11436-11443.
- Adamski, F. M., J. F. Atkins, and R. F. Gesteland. 1996. Ribosomal protein L9 interactions with 23 S rRNA: the use of a translational bypass assay to study the effect of amino acid substitutions. *J. Mol. Biol.* 261:357-371.
- Addinall, S. G., C. Cao, and J. Lutkenhaus. 1997. FtsN, a late recruit to the septum in *Escherichia coli*. *Mol. Microbiol.* 25:303-309.
- Addinall, S. G., and J. Lutkenhaus. 1996. FtsA is localized to the septum in an FtsZ-dependent manner. *J. Bacteriol.* 178:7167-7172.
- Adhya, S., P. P. Cleary, and A. Campbell. 1968. A deletion analysis of prophage lambda and adjacent genetic regions. *Proc. Natl. Acad. Sci. USA* 61:956-962.
- Adler, H. I., R. Mural, and B. Suttle. 1992. Oxygen sensitivity of an *Escherichia coli* mutant. *J. Bacteriol.* 174:2072-2077.
- Adler, H. I., W. D. Fisher, A. A. Hardigree, and G. E. Stapleton. 1966. Repair of radiation-induced damage to the cell division mechanism of *Escherichia coli*. *J. Bacteriol.* 91:737-742.
- Ahmad, S. I. 1996. A mutant of *Escherichia coli* hyper-resistant to a number of DNA damaging agents: location of the mutational site. *J. Photochem. Photobiol. B* 36:47-53.
- Ahmad, S. I., and R. H. Pritchard. 1969. A map of four genes specifying enzymes involved in catabolism of nucleosides and deoxynucleosides in *Escherichia coli*. *Mol. Gen. Genet.* 104:351-359.
- Ahmad, S. I., and R. H. Pritchard. 1972. Location of the gene specifying cytosine deaminase in *Escherichia coli*. *Mol. Gen. Genet.* 118:323-325.
- Ahmer, B. M., M. G. Thomas, R. A. Larsen, and K. Postle. 1995. Characterization of the *exbBD* operon of *Escherichia coli* and the role of ExbB and ExbD in TonB function and stability. *J. Bacteriol.* 177:4742-4747.
- Ahn, J., P. E. March, H. E. Takiff, and M. Inouye. 1986. A GTP-binding protein of *Escherichia coli* has homology to yeast RAS proteins. *Proc. Natl. Acad. Sci. USA* 83:8849-8853.
- Aiba, H. 1985. Transcription of the *Escherichia coli* adenylate cyclase gene is negatively regulated by cAMP-cAMP receptor protein. *J. Biol. Chem.* 260:3063-3070.
- Aiba, H., T. Baba, K. Hayashi, T. Inada, K. Isono, T. Itoh, H. Kasai, K. Kashimoto, S. Kimura, M. Kitakawa, M. Kitagawa, K. Makino, T. Miki, K. Mizobuchi, H. Mori, T. Mori, K. Motomura, S. Nakade, Y. Nakamura, H. Nishimoto, Y. Nishio, T. Oshima, N. Saito, G. Sampei, Y. Seki, S. Sivasundaram, H. Tagami, J. Takeda, K. Takemoto, Y. Takeuchi, C. Wada, Y. Yamamoto, and T. Horiuchi. 1996. GenBank submissions.
- Aiba, H., and K. Mizobuchi. 1989. Nucleotide sequence analysis of genes *purH* and *purD* involved in the *de novo* purine nucleotide biosynthesis of *Escherichia coli*. *J. Biol. Chem.* 264:21239-21246.
- Aiba, H., K. Mori, M. Tanaka, T. Ooi, A. M. Roy, and A. Danchin. 1984. The complete nucleotide sequence of the adenylate cyclase gene of *Escherichia coli*. *Nucleic Acids Res.* 12:9427-9440.
- Aiba, H., M. Kawamukai, and A. Ishihama. 1983. Cloning and promoter analysis of the *Escherichia coli* adenylate cyclase gene. *Nucleic Acids Res.* 11:3451-3465.
- Aiba, H., S. Adhya, and B. de Crombrughe. 1981. Evidence for two functional *gal* promoters in intact *Escherichia coli* cells. *J. Biol. Chem.* 256:11905-11910.
- Aiba, H., S. Fujimoto, and N. Ozaki. 1982. Molecular cloning and nucleotide sequencing of the gene for *E. coli* cAMP receptor protein. *Nucleic Acids Res.* 10:1345-1361.
- Aiba, H., T. Baba, K. Fujita, K. Hayashi, T. Inada, K. Isono, T. Itoh, H. Kasai, K. Kashimoto, S. Kimura, M. Kitakawa, K. Makino, T. Miki, K. Mizobuchi, H. Mori, T. Mori, K. Motomura, S. Nakade, Y. Nakamura, H. Nishimoto, Y. Nishio, T. Oshima, N. Saito, G. Sampei, Y. Seki, S. Sivasundaram, H. Tagami, J. Takeda, K. Takemoto, Y. Takeuchi, C. Wada, Y. Yamamoto, and T. Horiuchi. 1996. A 570-kb DNA sequence of the *Escherichia coli* K-12 genome corresponding to the 28.0-40.1 min region on the linkage map. *DNA Res.* 3:363-377.
- Aiso, T., and R. Ohki. 1998. An *meI* *pnp-7* double mutation suppresses the temperature-sensitive defect of *lacZ* gene expression in a *divE* mutant. *J. Bacteriol.* 180:1389-1395.
- Aizenman, E., H. Engelberg-Kulka, and G. Glaser. 1996. An *Escherichia coli* chromosomal "addiction module" regulated by 3',5'-bisphosphosphate: a model for programmed bacterial cell death. *Proc. Natl. Acad. Sci. USA* 93:6059-6063.
- Akamaru, J., S. Matsuyama, H. Tokuda, and S. Mizushima. 1991. Reconstitution of a protein translocation system containing purified SecY, SecE, and SecA from *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 88:6545-6549.
- Akiyama, M., E. Crooke, and A. Kornberg. 1992. The polyphosphate

- ported by the *Escherichia coli* maltose transport system. *J. Bacteriol.* 144: 7-11.
1225. Ferguson, G. P., A. D. Chacko, C. Lee, and I. R. Booth. 1996. The activity of the high-affinity K^+ uptake system Kdp sensitizes cells of *Escherichia coli* to methylglyoxal. *J. Bacteriol.* 178:3957-3961. (Erratum, 179:568, 1997.)
 1226. Ferguson, G. P., R. I. Creighton, Y. Nikolaev, and I. R. Booth. 1998. Importance of RpoS and Dps in survival of exposure of both exponential- and stationary-phase *Escherichia coli* cells to the electrophile *N*-ethylmaleimide. *J. Bacteriol.* 180:1030-1036.
 1227. Fernley, R. T., S. R. Lentz, and R. A. Bradshaw. 1981. Malate dehydrogenase isolation from *Escherichia coli* and comparison with the eukaryotic mitochondrial and cytoplasmic forms. *Biosci. Rep.* 1:497-507.
 1228. Ferone, R., S. C. Singer, M. H. Hanlon, and S. Roland. 1983. Isolation and characterization of an *E. coli* mutant affected in dihydrofolate- and folylpolyglutamate-synthetase, p. 585-589. In J. Blair (ed.), *Chemistry and biology of pteridines*. W. de Gruyter, New York, N.Y.
 1229. Fernandez, A., J. L. Garcia, and E. Diaz. 1997. Genetic characterization and expression in heterologous hosts of the 3-(3-hydroxyphenyl)propionate catabolic pathway of *Escherichia coli* K-12. *J. Bacteriol.* 179:2573-2581.
 1230. Fernandez, A., M. A. Prieto, J. L. Garcia, and E. Diaz. 1997. Molecular characterization of PadA, a phenylacetaldehyde dehydrogenase from *Escherichia coli*. *FEBS Lett.* 406:23-27.
 1231. Ferro-Novick, S., M. Honma, and J. R. Beckwith. 1984. The product of gene *secC* is involved in the synthesis of exported proteins in *E. coli*. *Cell* 38:211-217.
 1232. Feulner, G., J. A. Gray, J. A. Kirschman, A. F. Lehner, A. B. Sadosky, D. A. Vlazny, J. Zhang, S. Zhao, and C. W. Hill. 1990. Structure of the *rhsA* locus from *Escherichia coli* K-12 and comparison of *rhsA* with other members of the *rhs* multigene family. *J. Bacteriol.* 172:446-456.
 1233. Feutrier, J., M. Lepelletier, M. C. Pascal, and M. Chippaux. 1982. Tn10 insertions directed in the *pyrD-serC* region and improved mapping of *pepN* in *Escherichia coli* K12. *Mol. Gen. Genet.* 185:518-519.
 1234. Fiedler, W., and H. Rotering. 1985. Characterization of an *Escherichia coli* *mdoB* mutant strain unable to transfer *sn*-1-phosphoglycerol to membrane-derived oligosaccharides. *J. Biol. Chem.* 260:4799-4806.
 1235. Fiedler, W., and H. Rotering. 1988. Properties of *Escherichia coli* mutants lacking membrane-derived oligosaccharides. *J. Biol. Chem.* 263:14684-14689.
 1236. Figge, R. M., T. M. Ramseier, and M. H. Saier. 1994. The mannitol repressor (MtlR) of *Escherichia coli*. *J. Bacteriol.* 176:840-847.
 1237. Fill, N., and D. Friesen. 1968. Isolation of "relaxed" mutants of *Escherichia coli*. *J. Bacteriol.* 95:729-731.
 1238. Fill, N., J. D. Friesen, W. L. Downing, and P. P. Dennis. 1980. Post-transcriptional regulatory mutants in a ribosomal protein-RNA polymerase operon of *E. coli*. *Cell* 19:837-844.
 1239. Fijalkowska, I. J., R. L. Dunn, and R. M. Schaaper. 1997. Genetic requirements and mutational specificity of the *Escherichia coli* SOS mutator activity. *J. Bacteriol.* 179:7435-7445.
 1240. Fijalkowska, I. J., and R. M. Schaaper. 1995. Effects of *Escherichia coli* *dnaE* antimutator alleles in a proofreading-deficient *mutD5* strain. *J. Bacteriol.* 177:5979-5986.
 1241. Fikes, J. D., and P. J. Bassford. 1989. Novel *secA* alleles improve export of maltose-binding protein synthesized with a defective signal peptide. *J. Bacteriol.* 171:402-409.
 1242. Filpula, D. 1978. Regulation of ribonucleoside diphosphate reductase synthesis in *Escherichia coli*: specific activity of the enzyme in relationship to perturbations of DNA replication. *J. Bacteriol.* 135:429-435.
 1243. Filpula, D., and J. A. Fuchs. 1977. Regulation of ribonucleoside diphosphate reductase synthesis in *Escherichia coli*: increased enzyme synthesis as a result of inhibition of deoxyribonucleic acid synthesis. *J. Bacteriol.* 130:107-113.
 1244. Filutowicz, M., W. Ross, J. R. Wild, and R. L. Gourse. 1992. Involvement of Fis protein in replication of the *Escherichia coli* chromosome. *J. Bacteriol.* 174:398-407.
 1245. Fimmel, A. L., and B. A. Haddock. 1979. Use of *chlC-lac* fusions to determine regulation of gene *chlC* in *Escherichia coli* K-12. *J. Bacteriol.* 138:726-730.
 1246. Fimmel, A. L., and B. A. Haddock. 1981. Characterization of the *Escherichia coli* *chlC* regulatory region in a cloned *chlC-lac* gene fusion. *FEMS Microbiol. Lett.* 12:125-129.
 1247. Fimmel, A. L., and R. E. Loughlin. 1977. Isolation and characterization of *cysK* mutants of *Escherichia coli*. *J. Gen. Microbiol.* 103:37-43.
 1248. Finch, P. W., A. Storey, K. Brown, I. D. Hickson, and P. T. Emmerson. 1986. Complete nucleotide sequence of *recD*, the structural gene for the alpha subunit of exonuclease V of *Escherichia coli*. *Nucleic Acids Res.* 14:8583-8594.
 1249. Finch, P. W., A. Storey, K. E. Chapman, K. Brown, I. D. Hickson, and P. T. Emmerson. 1986. Complete nucleotide sequence of the *Escherichia coli* *recB* gene. *Nucleic Acids Res.* 14:8573-8582.
 1250. Finch, P. W., and P. T. Emmerson. 1983. Nucleotide sequence of the regulatory region of the *uvrD* gene of *Escherichia coli*. *Gene* 25:317-323.
 1251. Finch, P. W., and P. T. Emmerson. 1984. The nucleotide sequence of the *uvrD* gene of *E. coli*. *Nucleic Acids Res.* 12:5789-5799.
 1252. Finch, P. W., R. E. Wilson, K. Brown, I. D. Hickson, A. E. Tomlinson, and P. T. Emmerson. 1986. Complete nucleotide sequence of the *Escherichia coli* *recC* gene and of the *thyA-recC* intergenic region. *Nucleic Acids Res.* 14:4437-4451.
 1253. Finch, P. W., R. E. Wilson, K. Brown, I. D. Hickson, and P. T. Emmerson. 1986. Complete nucleotide sequence of *Escherichia coli* *ptr* gene encoding protease III. *Nucleic Acids Res.* 14:7695-7703.
 1254. Fischer, E., K. Gunter, and V. Braun. 1989. Involvement of ExbB and TonB in transport across the outer membrane of *Escherichia coli*: phenotypic complementation of *exb* mutants by overexpressed *tonB* and physical stabilization of TonB by ExbB. *J. Bacteriol.* 171:5127-5134.
 1255. Fischer, M., and S. A. Short. 1982. The cloning of the *Escherichia coli* K-12 deoxyribonucleoside operon. *Gene* 17:291-298.
 1256. Fischl, A. S., and E. P. Kennedy. 1990. Isolation and properties of acyl carrier protein phosphodiesterase of *Escherichia coli*. *J. Bacteriol.* 172: 5445-5449.
 1257. Flachmann, R., N. Kunz, J. Seifert, M. Gutlich, F. J. Wientjes, A. Laufer, and H. G. Gassen. 1988. Molecular biology of pyridine nucleotide biosynthesis in *Escherichia coli*. Cloning and characterization of quinolate synthesis genes *nadA* and *nadB*. *Eur. J. Biochem.* 175:221-228.
 1258. Flamm, E. L., and R. Weisberg. 1983. Primary structure of the *hip* gene of *Escherichia coli* and of its product, the β subunit of integration host factor. *J. Mol. Biol.* 183:117-128.
 1259. Flamm, J. A., J. D. Friesen, and A. J. Otsuka. 1988. The nucleotide sequence of the *Escherichia coli* *ns* gene. *Gene* 74:555-558.
 1260. Flannigan, K. A., S. H. Hennigan, H. H. Vogelbacker, J. S. Gots, and J. M. Smith. 1990. Purine biosynthesis in *Escherichia coli* K12: structure and DNA sequence studies of the *purHD* locus. *Mol. Microbiol.* 4:381-392.
 1261. Fleming, T. P., M. S. Nahlik, J. B. Neilands, and M. A. McIntosh. 1985. Physical and genetic characterization of cloned enterobactin genomic sequences from *Escherichia coli* K-12. *Gene* 34:47-54.
 1262. Fleming, T. P., M. S. Nahlik, and M. A. McIntosh. 1983. Regulation of enterobactin iron transport in *Escherichia coli*: characterization of *ent::Mu d(Ap^r lac)* operon fusions. *J. Bacteriol.* 156:1171-1177.
 1263. Flores, A., and J. Casades. 1995. Suppression of the pleiotropic effects of HisH and HisF overproduction identifies four novel loci on the *Salmonella typhimurium* chromosome: *osmH*, *sfiW*, *sfiX*, and *sfiY*. *J. Bacteriol.* 177:4841-4850.
 1264. Flower, A. M., and C. S. McHenry. 1986. The adjacent *dnaZ* and *dnaX* genes of *Escherichia coli* are contained within one continuous open reading frame. *Nucleic Acids Res.* 14:8091-8101.
 1265. Flower, A. M., and C. S. McHenry. 1990. The gamma subunit of DNA polymerase III holoenzyme of *Escherichia coli* is produced by ribosomal frameshifting. *Proc. Natl. Acad. Sci. USA* 87:3713-3717.
 1266. Flower, A. M., and C. S. McHenry. 1991. Transcriptional organization of the *Escherichia coli* *dnaX* gene. *J. Mol. Biol.* 220:649-658.
 1267. Flower, A. M., R. C. Doebele, and T. J. Silhavy. 1994. PrlA and PrlG suppressors reduce the requirement for signal sequence recognition. *J. Bacteriol.* 176:5607-5614.
 1268. Fogliano, M., and P. F. Schendel. 1981. Evidence for the inducibility of the *uvrB* operon. *Nature* 289:196-198.
 1269. Foglino, M., S. Gharbi, and A. Lazdunski. 1986. Nucleotide sequence of the *pepN* encoding aminopeptidase N of *Escherichia coli*. *Gene* 49:303-309.
 1270. Fognini-Lefebvre, N., J. C. Lazzarone, and R. Portaler. 1987. *tolA*, *tolB* and *excC*, three cistrons involved in the control of pleiotropic release of periplasmic proteins by *Escherichia coli* K12. *Mol. Gen. Genet.* 209:391-395.
 1271. Fong, S. T., J. Camakaris, and B. T. Lee. 1995. Molecular genetics of a chromosomal locus involved in copper tolerance in *Escherichia coli* K-12. *Mol. Microbiol.* 15:1127-1137.
 1272. Fonstein, M., T. Nikolskaya, D. Zaporozhets, Y. Nikolsky, S. T. Kulakauskas, and A. Mironov. 1994. Tn10-mediated inversions fuse uridine phosphorylase (*udp*) and rRNA genes of *Escherichia coli*. *J. Bacteriol.* 176: 2265-2271.
 1273. Fontecave, M., J. Coves, and J. L. Pierre. 1994. Ferric reductases or flavin reductases? *Biometals* 7:3-8.
 1274. Forchhammer, K., K. P. Rucknagel, and A. Bock. 1990. Purification and biochemical characterization of SELB, a translation factor involved in selenoprotein synthesis. *J. Biol. Chem.* 265:9346-9350.
 1275. Forchhammer, K., W. Leinfelder, and A. Bock. 1989. Identification of a novel translation factor necessary for the incorporation of selenocysteine into protein. *Nature* 342:453-456.
 1276. Forchhammer, K., W. Leinfelder, K. Boesmler, B. Veprek, and A. Bock. 1991. Selenocysteine synthase from *Escherichia coli*. Nucleotide sequence of the gene (*selA*) and purification of the protein. *J. Biol. Chem.* 266:6318-6323.
 1277. Forster, J. W., and P. Strike. 1985. Organization and control of the *Escherichia coli* *uvrC* gene. *Gene* 35:71-82.

3070. Murray, K. D., and H. Bremer. 1996. Control of *spoT*-dependent ppGpp synthesis and degradation in *Escherichia coli*. *J. Mol. Biol.* **259**:41-57.
3071. Murray, N. E., J. A. Gough, B. Suri, and T. A. Bickle. 1982. Structural homologies among type I restriction-modification systems. *EMBO J.* **1**:535-539.
3072. Murray, N. E., and W. S. Kelley. 1979. Characterization of *λ*polA transducing phages: effective expression of the *E. coli* *polA* gene. *Mol. Gen. Genet.* **175**:77-87.
3073. Muse, W. B., and R. A. Bender. 1998. The *nac* (nitrogen assimilation control) gene from *Escherichia coli*. *J. Bacteriol.* **180**:1166-1173.
3074. Muto, A., M. Sato, T. Tadaki, M. Fukushima, C. Ushida, and H. Himeno. 1996. Structure and function of 10Sa RNA: trans-translation system. *Biochimie* **78**:985-991.
3075. Mutoh, N., K. Inokuchi, and S. Mizushima. 1982. Amino acid sequence of the signal peptides of OmpF, a major outer membrane protein of *Escherichia coli*. *FEBS Lett.* **137**:171-174.
3076. Mutoh, N., and M. Simon. 1986. Nucleotide sequence corresponding to five chemotaxis genes in *Escherichia coli*. *J. Bacteriol.* **165**:161-166.
3077. Mutoh, N., T. Nagasawa, and S. Mizushima. 1981. Specialized transducing bacteriophage lambda carrying the structural gene for a major outer membrane matrix protein of *Escherichia coli* K-12. *J. Bacteriol.* **145**:1085-1090.
3078. Muzzin, O., E. A. Campbell, L. Xia, E. Severinova, S. A. Darst, and K. Severinov. 1998. Disruption of *Escherichia coli* *hepA*, an RNA polymerase-associated protein, causes UV sensitivity. *J. Biol. Chem.* **273**:15157-15161.
3079. Myler, P. J., G. M. Venkataraman, M. J. Lodes, and K. D. Stuart. 1994. A frequently amplified region in *Leishmania* contains a gene conserved in prokaryotes and eukaryotes. *Gene* **148**:187-193.
3080. Mytelka, D. S., and M. J. Chamberlin. 1996. *Escherichia coli* *fljAZY* operon. *J. Bacteriol.* **178**:24-34.
3081. Nagano, Y., R. Matsuno, and Y. Sasaki. 1991. An essential gene of *Escherichia coli* that has sequence similarity to a chloroplast gene of unknown function. *Mol. Gen. Genet.* **228**:62-64.
3082. Nagasawa, H., Y. Sakagami, A. Suzuki, H. Suzuki, H. Hara, and Y. Hirota. 1989. Determination of the cleavage site involved in C-terminal processing of penicillin-binding protein 3 of *Escherichia coli*. *J. Bacteriol.* **171**:5890-5893.
3083. Nagasawa, S., K. Ishige, and T. Mizuno. 1993. Novel members of the two-component signal transduction genes in *Escherichia coli*. *J. Biochem. (Tokyo)* **114**:350-357.
3084. Nagasawa, S., S. Tokishita, H. Aiba, and T. Mizuno. 1992. A novel sensor-regulator protein that belongs to the homologous family of signal-transduction proteins involved in adaptive responses in *Escherichia coli*. *Mol. Microbiol.* **6**:799-807.
3085. Nagase, T., S. Ishii, and F. Imamoto. 1988. Differential transcriptional control of the two *rRNA^{Met}* genes of *Escherichia coli* K-12. *Gene* **67**:49-57.
3086. Nagel, R., A. Chan, and E. Rosen. 1994. *riv* and *recG* genes and the induced precise excision of *Tn10* in *Escherichia coli*. *Mutat. Res.* **311**:103-109.
3087. Naggert, J., M. L. Narasimhan, L. DeVeaux, H. Cho, Z. I. Randhawa, J. E. Cronan, B. N. Green, and S. Smith. 1991. Cloning, sequencing, and characterization of *Escherichia coli* thioesterase II. *J. Biol. Chem.* **266**:11044-11050.
3088. Nagy, P. L., A. Marolewski, S. J. Benkovic, and H. Zalkin. 1995. Formyltetrahydrofolate hydrolase, a regulatory enzyme that functions to balance pools of tetrahydrofolate and one-carbon tetrahydrofolate adducts in *Escherichia coli*. *J. Bacteriol.* **177**:1292-1298.
3089. Nagy, P. L., G. M. McCorkle, and H. Zalkin. 1993. *purU*, a source of formate for *purT*-dependent phosphoribosyl-N-formylglycinamide synthesis. *J. Bacteriol.* **175**:7066-7073.
3090. Nahlik, M. S., T. J. Brickman, B. A. Ozenberger, and M. A. McIntosh. 1989. Nucleotide sequence and transcriptional organization of the *Escherichia coli* enterobactin biosynthesis cistrons *entB* and *entA*. *J. Bacteriol.* **171**:784-790.
3091. Nahlik, M. S., T. P. Fleming, and M. A. McIntosh. 1987. Cluster of genes controlling synthesis and activation of 2,3-dihydroxybenzoic acid in production of enterobactin in *Escherichia coli*. *J. Bacteriol.* **169**:4163-4170.
3092. Naito, S., T. Kitani, T. Ogawa, T. Okazaki, and H. Uchida. 1984. *Escherichia coli* mutants suppressing replication-defective mutations of the ColE1 plasmid. *Proc. Natl. Acad. Sci. USA* **81**:550-554.
3093. Nakabeppu, Y., H. Kondo, and M. Sekiguchi. 1984. Cloning and characterization of the *alkA* gene of *Escherichia coli* that encodes 3-methyladenine DNA glycosylase II. *J. Biol. Chem.* **259**:13723-13729.
3094. Nakabeppu, Y., H. Kondo, S. I. Kawabata, S. Iwanaga, and M. Sekiguchi. 1985. Purification and structure of the intact Ada regulatory protein of *Escherichia coli* K12. O⁶-methyl-guanine-DNA methyltransferase. *J. Biol. Chem.* **260**:7281-7288.
3095. Nakabeppu, Y., T. Miyata, H. Kondo, S. Iwanaga, and M. Sekiguchi. 1984. Structure and expression of the *alkA* gene of *Escherichia coli* involved in adaptive response to alkylating agents. *J. Biol. Chem.* **259**:13730-13736.
3096. Nakagawa, J., and M. Matsushashi. 1982. Molecular divergence of a major peptidoglycan synthetase with transglycosylase-transpeptidase activities in *Escherichia coli*—penicillin binding protein 1Bs. *Biochem. Biophys. Res. Commun.* **105**:1546-1553.
3097. Nakagawa, J., S. Tamaki, and M. Matsushashi. 1979. Purified penicillin binding proteins 1Bs from *Escherichia coli* membrane showing activities of both peptidoglycan polymerase and peptidoglycan crosslinking enzyme. *Agric. Biol. Chem.* **43**:1379-1380.
3098. Nakahigashi, K., and H. Inokuchi. 1990. Nucleotide sequence between the *fadB* and the *rnaA* operon from *Escherichia coli*. *Nucleic Acids Res.* **18**:6439.
3099. Nakahigashi, K., and H. Inokuchi. 1990. Nucleotide sequence of the *fadA* and *fadB* genes from *Escherichia coli*. *Nucleic Acids Res.* **18**:4937.
3100. Nakahigashi, K., K. Miyamoto, K. Nishimura, and H. Inokuchi. 1992. Isolation and characterization of a light-sensitive mutant of *Escherichia coli* K-12 with a mutation in a gene that is required for the biosynthesis of ubiquinone. *J. Bacteriol.* **174**:7352-7359.
3101. Nakahigashi, K., K. Nishimura, K. Miyamoto, and H. Inokuchi. 1991. Photosensitivity of a protoporphyrin-accumulating, light-sensitive mutant (*visA*) of *Escherichia coli* K-12. *Proc. Natl. Acad. Sci. USA* **88**:10520-10524.
3102. Nakajima, N., H. Ozeki, and Y. Shimura. 1981. Organization and structure of an *E. coli* tRNA operon containing seven tRNA genes. *Cell* **23**:239-249.
3103. Nakajima, N., H. Ozeki, and Y. Shimura. 1982. In vitro transcription of the *supB-E* tRNA operon of *Escherichia coli*. *J. Biol. Chem.* **257**:11113-11120.
3104. Nakamura, H. 1979. Novel acriflavine resistance genes, *acrC* and *acrD*, in *Escherichia coli* K-12. *J. Bacteriol.* **139**:8-12.
3105. Nakamura, H., H. Murakami, I. Yamato, and Y. Anraku. 1988. Nucleotide sequence of the *cybB* gene encoding cytochrome b561 in *Escherichia coli* K12. *Mol. Gen. Genet.* **212**:1-5.
3106. Nakamura, H., I. Yamato, Y. Anraku, L. Lemieux, and R. B. Gennis. 1990. Expression of *cyoA* and *cyoB* demonstrates that the CO-binding heme component of the *Escherichia coli* cytochrome *o* complex is in subunit I. *J. Biol. Chem.* **265**:11193-11197.
3107. Nakamura, H., M. Maruyama, M. Soma, J.-I. Kato, H. Suzuki, and Y. Hirota. 1983. On the process of cellular division in *Escherichia coli*: nucleotide sequence of the gene for penicillin-binding protein. *Mol. Gen. Genet.* **191**:1-9.
3108. Nakamura, H., and S. Mizusawa. 1985. In vivo evidence that the *nusA* and *infB* genes of *E. coli* are part of the same multi-gene operon which encodes at least four proteins. *EMBO J.* **4**:527-532.
3109. Nakamura, K., and M. Inouye. 1979. DNA sequence of the gene for the outer membrane lipoprotein of *E. coli*: an extremely AT-rich promoter. *Cell* **18**:1109-1117.
3110. Nakamura, K., R. M. Pirtle, I. L. Pirtle, K. Takeishi, and M. Inouye. 1980. Messenger ribonucleic acid of the lipoprotein of the *Escherichia coli* outer membrane. II. The complete nucleotide sequence. *J. Biol. Chem.* **255**:210-216.
3111. Nakamura, M., M. Yamada, Y. Hirota, K. Sugimoto, A. Oka, and M. Takamami. 1981. Nucleotide sequence of the *asnA* gene coding for asparagine synthetase of *E. coli* K-12. *Nucleic Acids Res.* **9**:4669-4676.
3112. Nakamura, S., M. Nakamura, T. Kojima, and H. Yoshida. 1989. *gyrA* and *gyrB* mutations in quinolone-resistant strains of *Escherichia coli*. *Antimicrob. Agents Chemother.* **33**:254-255.
3113. Nakamura, Y. 1980. Hybrid plasmid carrying *Escherichia coli* genes for primase (*dnaG*) and RNA polymerase sigma factor (*spoD*): gene organization and control of their expression. *Mol. Gen. Genet.* **178**:487-497.
3114. Nakamura, Y., T. Kurihara, H. Saito, and H. Uchida. 1979. Sigma subunit of *Escherichia coli* RNA polymerase affects the function of *AN* gene. *Proc. Natl. Acad. Sci. USA* **76**:4593-4597.
- 3114a. Nakanishi, A., T. Oshida, T. Matsushita, S. Imajoh-Ohmi, and T. Ohnuki. 1998. Identification of DNA gyrase inhibitor (GyrI) in *Escherichia coli*. *J. Biol. Chem.* **273**:19333-19338.
3115. Nakao, T., I. Yamato, and Y. Anraku. 1987. Nucleotide sequence of *putC*, the regulatory region for the *put* regulon of *Escherichia coli* K12. *Mol. Gen. Genet.* **210**:364-368.
3116. Nakao, T., I. Yamato, and Y. Anraku. 1987. Nucleotide sequence of *putP*, the proline carrier gene of *Escherichia coli* K12. *Mol. Gen. Genet.* **208**:70-75.
3117. Nakashima, K., A. Sugiura, K. Kanamaru, and T. Mizuno. 1993. Signal transduction between the two regulatory components involved in the regulation of the *kdpABC* operon in *Escherichia coli*: phosphorylation-dependent functioning of the positive regulator KdpE. *Mol. Microbiol.* **7**:109-116.
3118. Nakashima, K., A. Sugiura, and T. Mizuno. 1993. Functional reconstitution of the putative *Escherichia coli* osmosensor, KdpD, into liposomes. *J. Biochem. (Tokyo)* **114**:615-621.
3119. Nakashima, K., K. Kanamaru, T. Mizuno, and K. Horikoshi. 1996. A novel member of the *cspA* family of genes that is induced by cold shock in *Escherichia coli*. *J. Bacteriol.* **178**:2994-2997.
3120. Nakata, A., H. Shinagawa, and M. Amemura. 1982. Cloning of alkaline phosphatase isozyme gene (*iap*) of *Escherichia coli*. *Gene* **19**:313-319.
3121. Nakata, A., M. Amemura, and H. Shinagawa. 1984. Regulation of the phosphate regulon in *Escherichia coli* K-12: regulation of the negative